



**Preserve Descriptions of
Land Managed by the
Williamson County Conservation
Foundation under the
Williamson County Regional
Habitat Conservation Plan**

SWCA Project Number 31724.01

June 2017

SUBMITTED TO:

Gary Boyd
Williamson County Conservation Foundation
219 Perry Mayfield
Leander, Texas 78641

SUBMITTED BY:

SWCA Environmental Consultants
4407 Monterey Oaks Boulevard
Building 1, Suite 110
Austin, Texas 78749

**PRESERVE DESCRIPTIONS OF LAND MANAGED BY THE
WILLIAMSON COUNTY CONSERVATION FOUNDATION UNDER THE
WILLIAMSON COUNTY REGIONAL HABITAT CONSERVATION PLAN**

Prepared for

GARY BOYD
WILLIAMSON COUNTY CONSERVATION FOUNDATION
219 Perry Mayfield
Leander, Texas 78641

Prepared by

Stephen Van Kampen-Lewis
SWCA ENVIRONMENTAL CONSULTANTS
4407 Monterey Oaks Boulevard
Building 1, Suite 110
Austin, Texas 78749
www.swca.com

Kemble White
CAMBRIAN ENVIRONMENTAL
4422 Pack Saddle Pass, No. 204
Austin, Texas 78745

SWCA Project Number 31724.01

1 June 2017



As a licensed professional
geoscientist I attest that the contents
of this report are complete and
accurate to the best of my
knowledge.

Kemble White Ph.D., P.G.
Cambrian Environmental
Texas Geoscience Firm Registration Number 50484
kwhite@cambrianenvironmental.com
(512) 663-0156



This page intentionally left blank.

CONTENTS

PRESERVE SUMMARY	VII
1.0 INTRODUCTION.....	1
1.1 Karst Ecosystem Delineations.....	1
1.1.1 Karst Zones and Karst Fauna Regions.....	1
1.1.2 Karst Fauna Area Concept.....	3
2.0 PRESERVE DESCRIPTIONS	5
2.1 Twin Springs Karst Fauna Area.....	5
2.1.1 General Information.....	5
2.1.2 Hydrogeology	7
2.1.3 Caves of the Twin Springs KFA.....	9
2.2 Beck Preserve	12
2.2.1 Introduction.....	12
2.2.2 Hydrogeology	14
2.2.3 Caves of the Beck Preserve.....	16
2.3 Cobbs Cavern Karst Fauna Area	22
2.3.1 Introduction.....	22
2.3.2 Hydrogeology	25
2.3.3 Caves of the Cobbs Cavern KFA.....	26
2.4 Millennium Preserve.....	31
2.4.1 Introduction.....	31
2.4.2 Hydrogeology	34
2.4.3 Caves of the Millennium Preserve	35
2.5 Wilco Preserve.....	42
2.5.1 Introduction.....	42
2.5.2 Hydrogeology	44
2.5.3 Caves of the Wilco Preserve	44
2.6 Chaos Cave Preserve	50
2.6.1 Introduction.....	50
2.6.2 Hydrogeology	50
2.6.3 Caves of the Chaos Preserve.....	53
2.7 Big Oak Cave Preserve.....	57
2.7.1 Introduction.....	57
2.7.2 Hydrogeology	57
2.7.3 Caves of the Big Oak Cave Preserve	59
2.8 Priscilla's Well Karst Fauna Area.....	61
2.8.1 Introduction.....	61
2.8.2 Hydrogeology	63
2.8.3 Caves of the Priscilla's Well KFA.....	64
2.9 Woodland Park Cave Preserve.....	66
2.9.1 Introduction.....	66
2.9.2 Hydrogeology	66
2.9.3 Caves of the Woodland Park Cave Preserve.....	71

2.10 Karankawa Cave Karst Fauna Area.....	74
2.10.1 Introduction.....	74
2.10.2 Hydrogeology	76
2.10.3 Caves of the Karankawa Cave Karst Fauna Area	78
2.11 Coffin Cave Preserve	82
2.11.1 Introduction.....	82
2.11.2 Hydrogeology	84
2.11.3 Caves of the Coffin Cave Preserve	86
2.12 Beck Commons Preserve	88
2.12.1 Introduction.....	88
2.12.2 Hydrogeology	90
2.12.3 Caves of the Beck Commons Preserve	91
2.13 Shaman Cave Karst Fauna Area.....	93
2.13.1 Introduction.....	93
2.13.2 Hydrogeology	95
2.13.3 Caves of the Shaman Cave Karst Fauna Area	96
3.0 LITERATURE CITED	102

FIGURES

Figure 1.	Williamson County Conservation Foundation preserve system map.....	2
Figure 2.	Twin Springs Karst Fauna Area (KFA) location map.....	6
Figure 3.	Twin Springs KFA geologic cross-section.....	8
Figure 4.	Beck Preserve location map.	13
Figure 5.	Beck Preserve geology.	15
Figure 6.	Cobbs Cavern KFA map showing Cobbs Spring and documented karst features.	23
Figure 7.	Cobbs Ranch geology and idealized springshed hydrology for Cobbs Spring.	24
Figure 8.	Cobbs Spring cross sectional representation showing hydrologic mechanism behind spring flow.....	26
Figure 9.	Geologic cross section across Millennium and Wilco Preserves highlighting karst erosion by Honey Bear Creek.	32
Figure 10.	Millennium Preserve map showing documented caves.....	33
Figure 11.	Millennium Preserve showing geological cross section.....	35
Figure 12.	Wilco Preserve map showing documented karst features.	43
Figure 13.	Chaos Preserve location map.	51
Figure 14.	Geologic cross section of the Chaos Preserve.....	52
Figure 15.	Big Oak Cave preserve location map.	58
Figure 16.	Priscilla's Well KFA location map.	62
Figure 17.	Woodland Park location map.	67
Figure 18.	Cat Cave conservation area.	68
Figure 19.	Duckworth Bat Cave conservation area.	69
Figure 20.	Cat and Duckworth Bat Caves hydrogeologic setting with local springs.	70
Figure 21.	Karankawa Cave KFA location map.....	75
Figure 22.	Karankawa Cave KFA hydrogeology.	77
Figure 23.	Location map for the Coffin Cave Preserve.	83
Figure 24.	Coffin Cave is roughly formed in the Edwards Formation's upper half.	85
Figure 25.	Beck Commons Preserve location map.....	89
Figure 26.	Shaman Cave KFA location map.	94
Figure 27.	Geologic cross section of the Shaman Cave KFA.....	95

PHOTOGRAPHS

Photograph 1.	Entrance to Sunless City Cave shortly after discovery.	10
Photograph 2.	Beck Bat Cave entrance.	16
Photograph 3.	Beck Horse Cave entrance.	17
Photograph 4.	Beck Tex-2 Cave entrance.	18
Photograph 5.	Beck Salamander Cave entrance.	19
Photograph 6.	Photograph of the interior of Cobbs Cavern.	27
Photograph 7.	Bone Cave harvestman collected in Cobbs Cavern.	28
Photograph 8.	<i>Batrisesodes texanus</i> collected in Cobbs Cavern.	28
Photograph 9.	<i>Rhadine noctivaga</i> collected in Cobbs Cavern.	28
Photograph 10.	<i>Speodesmus bicornourus</i> collected in Cobbs Cavern.	28
Photograph 11.	<i>Cicurina vibora</i> collected in Cobbs Cavern.	29
Photograph 12.	<i>Eurycea</i> salamander at Cobbs Spring.	29
Photograph 13.	<i>Tayshaneta anopica</i> collected in Cobbs Cavern.	29
Photograph 14.	Through Trip Cave west entrance.	36
Photograph 15.	Little Demon Cave entrance.	37
Photograph 16.	Entrance to Millennium Cave.	38
Photograph 17.	Entrance to Fence and Trail Cave.	39
Photograph 18.	Entrance to Knuckle Cave.	40
Photograph 19.	Entrance to Wilco Cave.	46
Photograph 20.	Entrance to Choya Cave.	47
Photograph 21.	Entrance to Circulation Sink.	48
Photograph 22.	Chaos Cave entrance.	53
Photograph 23.	Under the Fence Cave entrance.	54
Photograph 24.	Poison Ivy Cave entrance.	55
Photograph 25.	Big Oak Cave western entrance.	59
Photograph 26.	Three visible best management practices at the Priscilla's Well KFA.	63
Photograph 27.	Cat Cave entrance.	71
Photograph 28.	Duckworth Bat Cave entrance.	72
Photograph 29.	Duckworth Bat Cave entrance pit.	72
Photograph 30.	The Coffin Cave entrance is a vertical shaft dropping into the Edwards Formation from the contact with the overlying Georgetown Formation.	85
Photograph 31.	Coffin Cave entrance.	87
Photograph 32.	Inscription from Eugene Beck dating to 1941.	90
Photograph 33.	Beck Sewer Cave's phreatic passage development in the lower level.	91
Photograph 34.	Biota survey crew preparing to enter Shaman Cave.	96
Photograph 35.	Biota survey crew (2016) looking for karst invertebrates within Shaman Cave.	97
Photograph 36.	Entrance to Powwow Cave.	98

TABLES

Table PS1.	Williamson County RHCP Preserve Resources	vii
Table PS2.	Current preserve acreage versus expected cave acreage as determined by the Williamson County Regional Habitat Conservation Plan.....	viii
Table PS3.	Established karst fauna areas and documented endangered karst invertebrate presence.	viii
Table 1.	Twin Springs KFA snapshot showing basic preserve information.	5
Table 2.	Species previously documented from the Twin Springs KFA	11
Table 3.	Beck Preserve snapshot showing basic preserve information.	12
Table 4.	Species previously documented from the Beck Preserve.....	20
Table 5.	Cobbs Cavern KFA snapshot showing basic preserve information.	22
Table 6.	Species previously documented from the Cobbs Cavern KFA.....	30
Table 7.	Millennium Preserve snapshot showing basic preserve information.	31
Table 8.	Species previously documented from the Millennium Preserve.	41
Table 9.	Wilco Preserve snapshot showing basic preserve information.	42
Table 10.	Species previously documented from the Wilco Preserve.	49
Table 11.	Chaos Cave Preserve snapshot showing basic preserve information.	50
Table 12.	Species previously documented from the Chaos Cave Preserve.....	55
Table 13.	Big Oak Cave Preserve snapshot showing basic preserve information.....	57
Table 14.	Species previously documented from the Big Oak Cave Preserve.....	60
Table 15.	Priscilla's Well KFA snapshot showing basic preserve information.....	61
Table 16.	Species previously documented from the Priscilla's Well KFA.....	65
Table 17.	Woodland Park Cave Preserve snapshot showing basic preserve information.	66
Table 18.	Species previously documented from the Woodland Park Preserve.	73
Table 19.	Karankawa Cave KFA snapshot showing basic preserve information.....	74
Table 20.	Species previously documented from the Karankawa Cave KFA.....	80
Table 21.	Coffin Cave Preserve snapshot showing basic preserve information.....	82
Table 22.	Species previously documented from the Coffin Cave Preserve.	86
Table 23.	Beck Commons Preserve snapshot showing basic preserve information.....	88
Table 24.	Species previously documented from the Beck Commons KFA.	92
Table 25.	Species previously documented from the Shaman Cave KFA.....	100

APPENDICES

- Appendix A. Cave Maps of the Twin Springs Preserve
- Appendix B. Cave Maps of the Beck Preserve
- Appendix C. Cave Maps of the Southwest Williamson County Regional Park (Millennium and Wilco Preserves)
- Appendix D. Cave Maps of the Chaos Preserve
- Appendix E. Cave Maps of the Priscilla's Well Karst Fauna Area
- Appendix F. Cave Maps of the Woodland Park Cave Preserve
- Appendix G. Cave Maps of the Karankawa Cave Karst Fauna Area
- Appendix H. Cave Map of the Coffin Cave Preserve
- Appendix I. Cave Map of the Beck Commons Preserve
- Appendix J. Cave Maps of the Shaman Cave Karst Fauna Area

PRESERVE SUMMARY

The 13 karst feature preserves currently administered by the Williamson County Conservation Foundation (WCCF) range in size from approximately 10 acres up to 172.2 acres and contain a wide variety of explored and unexplored karst habitat features. Table PS1 shows the number of karst features contained within each preserve, as well as any karst fauna area (KFA) status acreage. In sum, as of the end of 2016, the WCCF oversees the management of 57 named karst features within approximately 895.3 preserve acres. Designated KFAs cover approximately 530.1 acres.

Table PS1. Williamson County RHCP Preserve Resources

Preserve Name	Year Preserve Established	Number of Named Karst Features	Approximate Preserve Acreage	Approximate KFA Acreage	Listed Species Found In Preserve
Twin Springs KFA	2009	2	172.2	172.2	<i>Texella reyesi</i> , <i>Batrises texanus</i> *
Beck Preserve	2009	7	44.5	0.0	<i>Texella reyesi</i>
Cobbs Cavern KFA*	2009	1	163.0	163.0	<i>Texella reyesi</i> , <i>Batrises texanus</i>
Wilco Preserve	2002	13	152.5	0.0	<i>Texella reyesi</i>
Millennium Preserve	2002	6	74.4	0.0	<i>Texella reyesi</i>
Chaos Cave Preserve*	2002	4	30.0	0.0	<i>Texella reyesi</i>
Big Oak Preserve*	2003	1	10.0	0.0	<i>Rhadine persephone</i>
Priscilla's Well KFA	2011	2	51.5	51.5	<i>Texella reyesi</i> , <i>Batrises cryptotexanus</i>
Woodland Park Cave Preserve	2012	2	10.2	0.0	<i>Texella reyesi</i>
Karankawa Cave KFA	2013	8	61.7	61.7	<i>Texella reyesi</i> , <i>Batrises cryptotexanus</i>
Beck Commons Preserve	2014	2	4.2	0.0	<i>Texella reyesi</i>
Coffin Cave Preserve	2014	1	39.4	0.0	<i>Texella reyesi</i> , <i>Batrises texanus</i>
Shaman Cave KFA	2016	8	81.7	81.7	<i>Texella reyesi</i> , <i>Batrises cryptotexanus</i>
Total		57	895.3	530.1	

* Easement, not owned by Williamson County

The Williamson County Regional Habitat Conservation Plan (RHCP) Section 9.3.2 indicates that, by 2025, WCCF shall acquire 700 karst preserve acres through direct purchase or acquisition of perpetual conservation easements. That same section indicates the County planned to purchase 500 karst preserve acres outright and acquire 200 karst easement acres from private landowners. The WCCF currently administers approximately 895.3 karst preserve/easement acres and has thus exceeded the karst preserve acreage predicted for this point in time in the RHCP (SWCA et al. 2008). Table PS2 shows predicted karst preserve acreage at the RHCP's time of writing versus the current karst preserve acreage administered by the WCCF.

Table PS2. Current preserve acreage versus expected cave acreage as determined by the Williamson County Regional Habitat Conservation Plan.

Preserve	Minimum Cave Preserve Acreage Required by 2025	Current Preserve Holdings (2016)
Williamson County Owned	500	692.3
Conservation Easement	200	203
Total	700	895.3

The RHCP prescribes that the WCCF shall establish and administer nine to fifteen 40- to 90-acre KFAs to enhance efforts towards listed karst invertebrate recovery; three KFAs to be distributed within the three Williamson County KFRs (North Williamson County, Georgetown, McNeil/Round Rock) covered by the RHCP. As of 2016 WCCF administers five KFAs in the North Williamson County KFR for three karst invertebrates (*Texella reyesi*, *Batrisodes texanus*, *B. cryptotexanus*). Table PS3 lists current KFAs and their associated endangered karst invertebrates.

Table PS3. Established karst fauna areas and documented endangered karst invertebrate presence.

KFA	KFR	<i>Texella reyesi</i>	<i>Batrisodes texanus</i>	<i>Batrisodes cryptotexanus</i>
Cobbs Cavern	North Williamson County	√	√	
Twin Springs	North Williamson County	√	√	
Priscilla's Well	North Williamson County	√		√
Shaman Cave	North Williamson County	√		√
Karankawa Cave	North Williamson County	√		√

1.0 INTRODUCTION

Williamson County (the County) and the Williamson County Conservation Foundation (WCCF) were issued an incidental take permit by the U.S. Fish and Wildlife Service (USFWS) in October 2008 to authorize take of four endangered species arising from a variety of covered land development activities in properties voluntarily enrolled within the Williamson County Regional Habitat Conservation Plan (RHCP). The RHCP was prepared by the County with funding and technical assistance from the USFWS. The RHCP supports an incidental take permit that authorizes the *take* of the Bone Cave harvestman (*Texella reyesi*), the Coffin Cave mold beetle (*Batrisodes texanus*¹), the golden-cheeked warbler (*Setophaga chrysoparia*; GCWA), and the black-capped vireo (*Vireo atricapilla*; BCVI); collectively defined as the Covered Species. Take is defined as “to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct” (Chapter 16 United States Code Section 1532(19)). Authorized take of the Covered Species is mitigated primarily through the establishment and management of preserves providing habitat for the above-mentioned Covered Species plus a suite of other species occurring in the same habitat. The WCCF preserve system (Figure 1) is intended to sustain high-quality habitat for species already on the endangered species list and to proactively conserve habitat to preclude the need to list other species. These other species include, but are not limited to, 20 troglobitic cave invertebrates and four species of spring-adapted *Eurycea* salamanders. Covered and additional species addressed in the RHCP are discussed in the management plan for the preserves (Van Kampen-Lewis and White 2017). Sites documented herein are categorized either as preserves or as Karst Fauna Areas (KFAs), a categorization determined by criteria set forth by the USFWS (see RHCP Section 5.3.1.1 for further details). This document describes the preserve sites, their associated caves, and any habitat therein dedicated to birds and/or salamanders listed by the USFWS as threatened or endangered.

1.1 Karst Ecosystem Delineations

1.1.1 Karst Zones and Karst Fauna Regions

The USFWS commissioned a study in 1991 that attempted to determine the likelihood of various geological units in Williamson and Travis Counties to contain karst features with potential habitat for cave-dwelling invertebrates after listing several karst invertebrate species as endangered in 1988 (Veni and Associates 1992). These karst zones were subsequently updated by Veni and Martinez (2007). The study resulted in zone delineation based on lithology, cave distribution, cave fauna distribution, and geologic controls on cave development. The zones are delineated as follows:

- Zone 1 - contains endangered cave species
- Zone 2 - high probability of endangered or endemic cave fauna
- Zone 3 - low probability of endangered or endemic cave fauna
- Zone 4 - does not contain endangered or endemic cave fauna

The difference between Zones 1 and 2 is most likely an artifact of limited sampling. Zones 1 and 2 together reflect the potential distribution of cavernous rock exposed at the surface that may harbor karst invertebrates.

¹ Chandler and Reddell (2001) split the listed *Batrisodes texanus* into two species, *B. texanus* and *B. cryptotexanus*, but the USFWS (2009b) does not recognize the split. Species identified as *B. cryptotexanus* are known from 15 caves, all in Williamson County (Chandler and Reddell 2001; D.S. Chandler, personal communication to K. White, 2006). Both species are considered federally endangered and are protected under the Endangered Species Act. This report will reference each respective *Batrisodes* species by their taxonomically accepted Latin name.

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan




Figure 1. Williamson County Conservation Foundation preserve system map.

The study also discussed the overall karst geography of the Austin region, potential geologic and geographic barriers to karst invertebrate dispersal, and limits of species distribution. Veni and Associates (1992) originally proposed 10 karst geologic areas for Travis and Williamson Counties. USFWS (1994a) later modified the karst geologic areas into eight karst fauna regions (KFR) that serve as de facto recovery units for listed karst invertebrates: South Travis County, Rollingwood, Central Austin, and Jollyville in Travis County; and McNeil/Round Rock, Cedar Park, Georgetown, and North Williamson County in Williamson County.

Each delineated KFR was thought to be bound by geological and hydrological barriers to troglobitic species distribution as the concept was originally presented. However, subsequent studies have shown that the boundaries proposed for the KFRs do not always correspond to the known species' boundaries (Cokendolpher 2004; Elliott 2004; Paquin and Hedin 2004, 2005; White et al. 2001). In addition, some species previously thought to be restricted to one KFR have been collected over several contiguous KFRs (see Paquin and Hedin 2004 for *Cicurina madla*; Ubick and Briggs 1992 for *T. reyesi*).

1.1.2 Karst Fauna Area Concept

The KFA concept was first proposed in the *Recovery Plan for Endangered Karst Invertebrates in Travis and Williamson Counties, Texas* (USFWS 1994a). KFAs were envisioned as a unit of occupied habitat protected for the benefit of listed troglobitic species recovery. According to the Recovery Plan, a KFA is described as:

...an area known to support one or more locations of a listed species and is distinct in that it acts as a system that is separated from other karst fauna areas by geologic and hydrologic features and/or processes that create barriers to the movement of water, contaminants, and troglobitic fauna (USFWS 1994a:76).

To be considered "protected", a karst fauna area should contain a large enough expanse of contiguous karst and surface area to maintain the integrity of the karst ecosystem on which each species depends. The size and configuration of each karst fauna area should be adequate to maintain moist, humid conditions, air flow, and stable temperatures in the air-filled voids; maintain an adequate nutrient supply; prevent contamination of surface and groundwater entering the ecosystem; prevent or control the invasion of exotic species, such as red imported fire ants; and allow for movement of the karst fauna and nutrients through the interstitial spaces between karst features (USFWS 1994a:82).

Although the Recovery Plan indicates that three KFAs should be set aside for perpetual protection within each KFR for each listed species, it provides only the following general guidelines for determining what conditions constitute a KFA:

Karst fauna areas should be far enough apart so that if a catastrophic event (for example, contamination of the water supply, flooding, disease) were to destroy one of the areas and/or the species in it, that event would not likely destroy any other area occupied by that species. (USFWS 1994a:76)

Other factors to consider when selecting KFAs include:

...the ability to ensure long-term protection, current level of habitat disturbance, past and present land use, presence of other rare or candidate species, ease of protection (landowner cooperation), and, where applicable, importance to the regional groundwater system. (USFWS 1994a:80).

...the pattern and direction of groundwater movement, direction and area of surface and subsurface drainage, preservation of the surface community above and surrounding the cave, and the presence of other caves or karst features. In general, land bounded by the contour interval at the cave floor is the area within which contaminants moving over the surface or through karst could move toward the cave. Outside this contour, contaminants would move away from the cave. (USFWS 1994a:82).

No specific criteria were provided in the Recovery Plan beyond these general guidelines. However, the RHCP indicates that minimum size for KFAs will be 40 acres where possible in order to protect the natural surface vegetation around a cave or cave cluster needed to support the cave ecosystem over the long term.

2.0 PRESERVE DESCRIPTIONS

2.1 Twin Springs Karst Fauna Area

2.1.1 General Information

Table 1. Twin Springs KFA snapshot showing basic preserve information.

Twin Springs KFA Snapshot	
Preserve Inception Year	Acquired parcels between 2007-2009
Acreage	172.2
Fence Status	Fully fenced, with repairs needed between U.S. Army Corps of Engineers land
Sign Status	Signs posted around perimeter, warning signs on all cave gates
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Sunless City, Whitney West
Non-gated Caves	None

The Twin Springs KFA includes 172.2 acres located adjacent to U.S. Army Corps of Engineers (USACE) land on the north side of Lake Georgetown, west of Russell Park Road and south of the end of Twin Springs Road (Figure 2). It comprises three contiguous management areas acquired by the County between 2007 and 2009. The 12.2-acre Sunless City Cave Preserve was transferred to the County by the Texas Department of Transportation (TxDOT) in 2007 as a mitigation parcel established for the State Highway (SH) 45 project. The County acquired the adjacent 145 acres in 2008 that were not developed under the terms of the Russell Park Estates Environmental Assessment (EA)/Habitat Conservation Plan (HCP). The County also acquired management responsibility for an additional 15 acres of deed-restricted open space within the Russell Park Estates development through a 2009 agreement established between the County and the Russell Park Neighborhood Association. These three adjacent areas are managed collectively as the 172.2-acre Twin Springs KFA; with no management differences between the three tracts. Access points include gates adjacent to County Road (CR) 262 and at the cul-de-sac at the end of Twin Springs Road.

The Twin Springs KFA is completely fenced and maintains appropriate signage to aid in trespassing prevention. However, the fence abutting the USACE land to the south of Twin Springs KFA is not in good shape; though trespassing from this location is unlikely to occur due to the neighboring tract's remote nature. Few fence gaps currently exist on the southern property line; however, a planned development along this area has prompted that these gaps be fixed.

A Leave No Trace (LNT) awareness course with specific adaptations for Williamson County preserves is taught for those who wish to access the Twin Springs KFA. This 2-hour-long course provides the site-specific plan for the Twin Springs KFA and an overview of the LNT program. Attendees are provided a permit to Twin Springs KFA (and other Williamson County preserves) after completing the LNT coursework. The permit must be worn at all times while inside the KFA and human impacts are currently being monitored. No detectable impact has been documented from visitor use. The WCCF issued 467 LNT passes in 2014, 113 passes in 2015, and 57 passes in 2016.




Figure 2. Twin Springs Karst Fauna Area (KFA) location map.

2.1.2 Hydrogeology

The Twin Springs KFA topography slopes from uplands in the northeast and northwest toward the south-central portion of the property to Taylor Ray Hollow, which drains to Lake Georgetown. Total relief on the Twin Springs KFA is approximately 140 feet, ranging from approximately 990 to 850 feet above mean sea level (amsl) (Figure 3). The Edwards Limestone is approximately 110 feet thick in the area of Twin Springs KFA. Local geology is almost entirely underlain by the Edwards limestone with the exception of the spring run and the lowest elevations along the southern preserve boundary, where the less-permeable Comanche Peak formation is exposed. The contact between these units, found generally at 880 feet amsl, provides a discharge point for the Edwards Aquifer known as Twin Springs (see Figure 3). However, recharge features on the property probably do not contribute water to the main body of the Edwards Aquifer because the hydraulic gradient between the uplands and the canyon bottom is far greater than that between the uplands and the base of the aquifer. Accordingly, recharging waters are more likely to discharge through seeps and springs near the contact between the Edwards and Comanche Peak formations (Jones 2003) as evidenced by discharge features within Twin Springs KFA.

The permanently flowing spring located in the western branch of Taylor Ray Hollow provides excellent habitat for the Georgetown salamander. Although discharge is variable, the spring has not been known to go dry even in the 2008 and 2009 record-breaking drought. This indicates substantial interconnectivity with the main body of the Edwards Aquifer and a relatively large springshed area extending well beyond Twin Springs KFA boundaries. Based on the geomorphology of the local portion of the San Gabriel River drainage basin and the known locations of other springs in adjacent tributaries, the springshed may extend more than a mile to the northeast to Ranch-to-Market Road (RM) 2338, where the Georgetown limestone covers Edwards limestone along the drainage divide between the North Fork San Gabriel River and Berry Creek. Development in this area consists primarily of single-family homes on 2- to 5-acre lots. The relatively low impervious cover associated with this development style may have contributed to maintaining spring flow quality and quantity.

Groundwater conduits within the drainage network tend to develop greater degrees of connectivity and transport efficiency in response to the hydraulic gradient between the land surface and the base level as a karst system matures. Base level is roughly equivalent to the water table in porous media aquifers. Conduits in the vadose zone (above the saturated zone) tend to develop increasingly vertical morphologies as this occurs in response to the steep-to-vertical hydraulic gradient between the land surface and the base level. Conduits reaching base level tend to follow horizontal flow paths along the relatively shallow, commonly nearly flat, hydraulic gradient between the base level and the resurgence through which groundwater discharges at the surface. As Sunless City Cave and Whitney West Cave are part of a relatively mature karst system, it is likely that recharging waters travel along predominantly vertical pathways until encountering horizontal flow paths at the Edwards/Comanche Peak contact where they are routed toward springs above the North Fork San Gabriel River. Accordingly, recharge entering the caves likely originates primarily from strata directly above the cave footprint and secondarily through lateral seepage from a relatively restricted perimeter around the footprint. It appears, therefore, that events occurring outside of the KFA pose relatively little contamination risk for KFA caves.




Figure 3. Twin Springs KFA geologic cross-section.

2.1.3 Caves of the Twin Springs KFA

Cave maps are included within Appendix A.

The Twin Springs KFA contains high-quality habitat for GCWA, Georgetown salamander (*Eurycea naufragia*), and two listed karst invertebrates. Twin Springs KFA value is additionally enhanced by adjacency to undeveloped habitat around Lake Georgetown. The USACE manages the natural resources around Lake Georgetown in cooperation with Texas Parks and Wildlife Department (TPWD), the USFWS, and the Texas Forest Service. Two known caves (Whitney West and Sunless City) and several unexcavated karst features occur within the Twin Springs KFA. At least eight troglobitic species occur within the KFA including two federally endangered species: the Bone Cave harvestman and *Batrissodes texanus* (Table 2). Both caves are confirmed locations for the Bone Cave harvestman and *Batrissodes texanus* has been confirmed from Sunless City Cave and likely occurs in both caves. The WCCF assumes that Twin Springs KFA will be recognized as contributing to *Batrissodes texanus* long-term recovery if that species is detected within Whitney West Cave.

Sunless City Cave: Sunless City Cave was discovered during the planning process for the existing development along Whitney Woods Circle at the eastern preserve boundary. During excavation at the base of a nearly perfectly circular solution sinkhole, a vertical shaft was discovered dropping approximately 20 feet into a single open chamber (Photograph 1). Sparkling white speleothems on the opposite side of the main room from the entrance shaft had the appearance of a nighttime city skyline and inspired the name Sunless City (a play on words referencing the nearby Sun City Development). The stratigraphic location of the cave at the base of the Edwards Aquifer and the general morphology of the cave, including phreatic pressure doming preserved in the ceiling, indicate that Sunless City Cave was formed under phreatic conditions (below the water table) prior to the draining of the local portion of the northern segment of the Edwards Aquifer by the down-cutting of the San Gabriel River. Slowly circulating ground water gradually dissolved and removed relatively soluble limestone in the strata in which the cave is formed, creating void space. This most likely occurred preferentially along rock joints and other fractures which are currently visible in the cave walls and ceilings. As the water table dropped due to stream incision and general denudation of the land surface, Sunless City Cave became perched above the water table in the vadose zone. Subsequent collapse and modification by vertically infiltrating vadose waters have contributed to the current morphology of the cave and largely obscured additional clues as to the genesis of the cave. The current *high and dry* nature of caves within Twin Springs KFA is what causes them to be high-quality habitat for a diverse community of air-breathing terrestrial karst invertebrates. This cave is known to contain the Bone Cave harvestman and *Batrissodes texanus*. A cave gate currently prevents unauthorized access to this feature.



Photograph 1. Entrance to Sunless City Cave shortly after discovery.

Whitney West Cave: Whitney West Cave was discovered after a lengthy and difficult excavation guided by geophysical survey in the summer of 2008. A 25-foot-wide, but relatively shallow, collapsed sinkhole ringed by large plateau live oaks (*Quercus fusiformis*) indicated the presence of sub-surface void space, but the size of breakdown slabs within the sinkhole interior and the lack of an obvious drain left little clue as to which direction to focus excavation efforts. An electrical resistivity survey of the sinkhole indicated that void space was present and that the southern rim of the sink was most proximal to the void. Investigators discovered a fracture-controlled conduit that consistently blew cool air following approximately one month of excavation through dense fractured limestone, thick root mats, and soil. That conduit was widened into a crawlspace large enough to conduct a biological survey. After repeated efforts, the Bone Cave harvestman was discovered along with several other troglobitic species. The conduit still blows air from the great cave system at the far end of the crawl space. The crawlspace appears to be located above and adjacent to a much larger void space of unknown dimensions according to the geophysical survey profile. It is presumed that Whitney West Cave will be as biologically diverse as Sunless City Cave if it is decided to continue excavation efforts in order to allow access to the main cavern for biological surveys. A cave gate currently prevents unauthorized access to Whitney West Cave.

Table 2 shows documented species within the Twin Springs KFA.

Table 2. Species previously documented from the Twin Springs KFA

Species		Sunless City Cave	Whitney West Cave
Permitted Species			
<i>Texella reyesi</i>		√	√
<i>Batrises texanus</i>		√	
Other Species			
Crickets	<i>Ceuthophilus cunicularis</i>	√	√
	<i>Ceuthophilus secretus</i>	√	√
	<i>Ceuthophilus</i> sp. B	√	√
Arachnids	<i>Cicurina varians</i>	√	√
	<i>Cicurina vibora</i>	√	√
	<i>Tayshaneta</i> sp.		√
	<i>Achaearanea</i> sp.		√
	<i>Eidmanella</i> sp.	√	√
	<i>Leiobunum</i> sp.		√
	<i>Pseudouroctonus reddelli</i>	√	√
Millipedes/ Centipedes	<i>Speodesmus bicornourus</i>	√	√
	<i>Cambala speobia</i>	√	
	<i>Scolopendra</i> sp.	√	
	Unknown Centipede	√	
	Scutigerae (House Centipede)	√	
Beetles	<i>Rhadine subterranea</i>	√	
	Staphylinidae	√	
	Coleoptera		√
Reptiles/ Amphibians	<i>Eleutherodactylus marnockii</i>	√	√
	<i>Incilius valliceps</i>	√	
Mammals	<i>Perimyotis subflavus</i>	√	
	<i>Myotis velifer</i>	√	
Other	Desert Cockroach	√	√
	Surface Isopod		√
	<i>Brackenridgia</i> sp., cf <i>cavernarum</i>	√	
	<i>Texoreddellia texensis</i>	√	
	<i>Collembola</i> sp.	√	√
	<i>Heliodiscus eigenmanni</i>	√	
	Earthworm	√	
	Assassin Bug		√
	Fly	√	
	Gnats		√

2.2 Beck Preserve

2.2.1 Introduction

Table 1. Beck Preserve snapshot showing basic preserve information.

Beck Preserve Snapshot	
Preserve Inception Year	2006
Acreage	44.5
Fence Status	Fence separates preserve from adjacent school. No other fence.
Sign Status	Warning signs on all cave gates
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Beck Tex-2, Beck Salamander, Beck Horse, Beck Bat, Beck Pride, Beck Creek and Beck Crevice Caves
Non-gated Caves	None

The Beck Preserve is approximately 44.5 acres purchased by the WCCF with funds from an Endangered Species Act (ESA) Section 6 grant provided by the USFWS and administered by the TPWD. It is located in the City of Round Rock, southwest from Ranch-to-Market Road (RM) 620 and Great Oaks Drive intersection (Figure 4). Beck Preserve is bounded to the west and south by Cedar Valley Middle School, to the north by Great Oaks Drive, and to the east by RM 620. The site includes seven caves and some have been studied for many years, including notable caves on the former Beck Ranch. Some of the first formal, scientific studies of Texas cave life were conducted on the Beck Ranch in the 1970s by Dr. Robert Mitchell of Texas Tech University. His studies indicate Beck Bat Cave, Beck Crevice Cave (which is now considered smaller feature within Beck Bat Cave), Beck Horse Cave, Beck Pride Cave, and Beck Tex-2 Cave are known to contain the Bone Cave harvestman. Two other caves, Beck Salamander Cave and Beck Creek Cave, are not known to contain endangered fauna but may be connected to the other caves. Each cave is considered an aquifer recharge feature. Beck Preserve may constitute a KFA, but has yet to be designated as such by the USFWS.




Figure 4. Beck Preserve location map.

The Beck Preserve is located at the southern end of one of the densest cave concentrations known within the Bone Cave harvestman range. Approximately 50 caves have been documented from an area measuring roughly 1.5 miles long north/south by 1 mile wide east/west. The Beck Preserve is adjacent to the Brushy Creek Municipal Utility District, which contains 13 additional caves on four nearly contiguous parcels beginning on the north side of Great Oaks Drive and stretching for a mile to the north beyond O'Connor Drive. Eight of those caves are confirmed to contain the Bone Cave harvestman. Approximately 500 feet northeast of the Beck Preserve between Great Oaks Drive and Scott and White Drive is a roughly 2-acre setback containing Vault and Imprint caves. Approximately 1,500 feet northeast of the Beck Preserve is a parcel containing Beck Sewer Cave and Beck Tin Can Cave maintained by the WCCF as the Beck Commons Preserve (Section 3.12). Immediately across RM 620 from Beck Preserve is the Highland Horizon tract which contains 12 caves occupied by the Bone Cave harvestman plus several additional caves. Robinson Ranch contains several occupied caves approximately 0.5 mile east from Beck Preserve, and additional acreage on the ranch has not been surveyed for karst features. Some of these caves are potentially connected to caves within Beck Preserve, at least from an invertebrate's point of view. Although this potential is difficult to quantify, it appears to be greatest at the following locations: Jackhammer Cave,² which is located partially under RM 620 approximately 275 feet from Beck Tex-2 Cave; Trouser Press Cave, which is located adjacent to RM 620 approximately 490 feet from Beck Tex-2 Cave; and Anteater's Delight³ Cave, which is located immediately adjacent to the east side of RM 620 less than 200 feet from the edge of the footprint of Beck Bat Cave.

The Beck Preserve currently maintains a fence separating it from the adjacent Cedar Valley Middle School. All other portions remain unfenced; however, staff from the middle school maintain a constant presence to prevent truant students from bypassing the existing fence and accessing Beck Preserve. Additionally, Beck Tex-2, Beck Salamander, Beck Horse, Beck Bat, Beck Pride and Beck Crevice Caves all have cave gates blocking access to their respective caves.

2.2.2 Hydrogeology

The Beck Preserve is located entirely within the Edwards Aquifer Recharge Zone (EARZ) (Figure 5). Based on regional trends, the direction of groundwater flow from Beck Preserve is likely the northeast. Recharge from Beck Preserve likely contributes baseflow to Brushy Creek, but no formal study has been conducted.

² Jackhammer Cave was sealed as result of RM 620 expansion and was a confirmed *Texella reyesi* locality.

³ Anteater's Delight was sealed as a result of development activity and was not a known *Texella reyesi* locality.





Figure 5. Beck Preserve geology.

2.2.3 Caves of the Beck Preserve

Cave maps are included within Appendix B.

Beck Bat Cave: Beck Bat Cave is the most notable feature on the site with records kept by cave explorers as early as the 1950s (Photograph 2). Beck Bat Cave contained a colony of “Mexican brown bats” (*Myotis velifer*, now known as cave myotis) as recently as 1989, but disturbance from human visitation prior to preserve acquisition and cave gate installation caused them to relocate. The cave entrance measures approximately 8 feet in diameter, sloping to the north beneath a ledge of bedrock. The cave primarily consists of a single large, wide room, divided into two main areas by a sizable ceiling collapse. Two small, lower-level rooms are at the northern extent of the cave. The cave dimensions measure approximately 163 feet long by 91 feet wide, with a maximum depth of 22 feet. This cave is known to contain the Bone Cave harvestman (Table 4). A cave gate currently prevents unauthorized access to this feature.



Photograph 2. Beck Bat Cave entrance.

Beck Crevice Cave: Beck Crevice Cave is approximately 100 feet from the entrance of Beck Bat Cave and two features share structural connection. The cave entrance drops 5 feet into a low bedding plane room that extends for approximately 30 feet towards Beck Bat Cave. The passage is formed along a rock joint trending approximately N40E. The footprint of the cave measures approximately 22 feet wide by 38 feet long, with a maximum depth of 18.2 feet. This cave is known to contain the Bone Cave harvestman (see Table 4) but is no longer monitored due to its small size and connectivity with the larger, Beck Bat Cave.

Beck Horse Cave: The Beck Horse Cave entrance is approximately 10 feet in diameter, sloping beneath the north headwall into a large room (Photograph 3). The cave is formed along a rock joint trending

N21E. This cave is known to contain the Bone Cave harvestman (see Table 4). A cave gate currently prevents unauthorized access to this feature.



Photograph 3. Beck Horse Cave entrance.

Beck Pride Cave: Beck Pride Cave is Beck Preserve's most biologically diverse cave. The cave entrance drops approximately 15 feet to the main room, which was formed along a rock joint trending approximately N16E. The cave contains upper-, mid-, and lower-level extents, all controlled along this same rock joint. The main passage extends to the north mainly as a walking size passage until it pinches off at a low bedding plane. The cave also extends to the south along the mid-level as a low, wide bedding plane passage. This cave is known to contain the Bone Cave harvestman (see Table 4). A cave gate currently prevents unauthorized access to this feature.

Beck Tex-2 Cave: Beck Tex-2 Cave is very shallowly developed, with a bedding plane level beginning at 3 feet below the ground surface that slopes down to the west (Photograph 4). The surveyed length of the cave is 69.6 feet. The cave footprint measures approximately 20 feet wide by 32 feet long, with a maximum depth of 10.6 feet. This cave is known to contain the Bone Cave harvestman (see Table 4). A cave gate currently prevents unauthorized access to this feature.



Photograph 4. Beck Tex-2 Cave entrance.

Beck Salamander Cave: The Beck Salamander Cave entrance drops vertically into an enlarged rock joint trending N30E (Photograph 5). The cave measures approximately 11.4 feet long, with a maximum depth of 12.1 feet. This cave is not currently known to contain any endangered karst invertebrates (see Table 4). A cave gate currently prevents unauthorized access to this feature.



Photograph 5. Beck Salamander Cave entrance.

Beck Creek Cave: The Beck Creek Cave entrance headwall resembles that of Beck Bat Cave and Beck Horse Cave; however, the passage is completely blocked by massive in-filled rocks and sediments. The cave was backfilled with the use of a dozer in the past, and an attempt to re-open the cave was unsuccessful. This cave is not currently known to contain any endangered karst invertebrates (see Table 4). A cave gate currently prevents unauthorized access to this feature, though this cave is not currently surveyed for karst biota.

Table 4 shows documented species within the Beck Preserve.

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan

Table 2. Species previously documented from the Beck Preserve.

Species		Beck Bat Cave	Beck Pride Cave	Beck Horse Cave	Beck Crevice Cave	Beck Tex-2 Cave	Beck Salamander Cave	Beck Creek Cave
Permitted Species								
<i>Texella reyesi</i>		√	√	√	√	√		
Other Species								
Crickets	<i>Ceuthophilus</i> sp.	√	√	√	√	√	√	√
	<i>Cicurina varians</i>	√	√	√	√	√		
Arachnids	<i>Cicurina buwata</i>	√	√		√	√		
	<i>Tayshaneta</i> sp.	√						
	<i>Eperigone albula</i>	√						
	<i>Anapistula</i> sp.	√		√				
	<i>Agyneta llanoensis</i>	√						
	Araneae	√	√	√	√	√	√	√
	<i>Eidmannella</i> sp.	√		√				
	<i>Achaeareanea</i> sp.				√	√		
	<i>Pseudouroctonus reddelli</i>	√						
	<i>Lechytia</i> sp.			√				
	<i>Tartarocreagris infernalis</i>	√	√					
Millipedes/ Centipedes	<i>Speodesmus bicornourus</i>	√	√	√			√	
	<i>Cambala speobia</i>	√	√	√	√	√		
	<i>Oxidus gracilis</i>			√	√			
	<i>Myrmecodesmus</i> sp.		√					
	<i>Scolopendra</i> sp.		√				√	
	Scutigeridae		√			√		
Beetles	<i>Rhadine subterranea</i>	√	√					
	Coleoptera	√			√		√	
	Staphylinidae	√		√	√		√	√
	<i>Eustilicus condei</i>	√						
	<i>Biblioplectus</i> sp. nr. <i>ruficeps</i>			√				
	<i>Batrisodes uncicornis</i>	√	√	√	√	√		
	<i>Tachys ferrugineus</i>	√						
	Tachyini							√
Reptiles/ Amphibians	<i>Plethodon albagula</i>						√	
	<i>Eleutherodactylus marnockii</i>							

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan

Species		Beck Bat Cave	Beck Pride Cave	Beck Horse Cave	Beck Crevice Cave	Beck Tex-2 Cave	Beck Salamander Cave	Beck Creek Cave
Mammals	<i>Perimyotis subflavus</i>	√	√	√				
	<i>Myotis velifer</i>	√		√				
	<i>Procyon lotor</i>		√					
Other	Desert Cockroach		√					
	Blattaria							√
	<i>Collembola</i> sp.	√	√	√	√	√	√	√
	Isopod (pillbug)	√	√	√	√	√		√
	Earthworm							√
	Gastropoda (Slugs/Snails)	√	√	√		√	√	√
	Formicidae			√	√			
	<i>Solenopsis invicta</i>	√	√	√		√	√	√
	Lepidoptera	√						
	Psocoptera			√				
	Thysanoptera			√				
	Hymenoptera					√		
	Flea	√					√	
	Fly			√		√	√	
	Gnat	√					√	
	Mite	√	√		√			√
	Mosquito						√	
	Streblidae	√						
	Diptera	√		√				
	<i>Helicodiscus eigenmanni</i>	√		√		√	√	
	<i>Coragyps atratus</i>							

2.3 Cobbs Cavern Karst Fauna Area

2.3.1 Introduction

Table 3. Cobbs Cavern KFA snapshot showing basic preserve information.

Cobbs Cavern KFA Snapshot	
Preserve Inception Year	Acquired easements 2006, 2011
Acreage	163.0
Fence Status	None- However, the greater ranch is fenced
Sign Status	None- Private property
Baseline Vegetation Survey Date	None
Owner	Easement- Lyda Family Trust
Gated Caves	Cobbs Cavern
Non-gated Caves	None

The Cobbs Ranch comprises approximately 1,670 acres and is approximately 5 miles northwest from the City of Georgetown (Figure 6). The WCCF received a Recovery Land Acquisition grant through the Cooperative Endangered Species Conservation Fund and authorized by Section 6 of the ESA, to provide 75% of the cost to purchase a conservation easement. The federal grant required a 25% local match, which was donated by the Lyda Family Trust as land. The grant was awarded to the County in September 2005. The County executed a conservation easement contract with the Lyda Family Trust to establish the 64.4-acre conservation easement in November 2006. The WCCF was awarded additional Section 6 funds in 2011, which were used to acquire the conservation easement for an additional 98 acres surrounding the entire Cobbs Cavern footprint. This acquisition completed the approximately 163.0-acre Cobbs Cavern KFA, which was approved by the USFWS in October 2011. The ranch consists of live oak-juniper savanna on gently rolling uplands drained generally to the east by the Cobbs Spring branch and other tributaries to Berry Creek. The property is located within the northern Balcones Fault Zone (BFZ) and within the northern segment of the EARZ.

The spring system that created the extensive cavern system created a multitude of ecological niches inhabited by some of the rarest, most unique, and endangered organisms in Texas. Cobbs Cavern is one of the 30 longest caves in the state, with over 4,500 feet of passage, all of which is located within the conservation easement. It is home to at least six species of rare and endangered terrestrial karst invertebrates, including two covered species and four additional species as listed in the RHCP (Table 6). The conservation easement contains a significant portion of the springshed for Cobbs Spring, which is known to contain a population of the Georgetown salamander (Figure 7).




Figure 6. Cobbs Cavern KFA map showing Cobbs Spring and documented karst features.




Figure 7. Cobbs Ranch geology and idealized springshed hydrology for Cobbs Spring.

Certain activities were mandated by the final conditions of the Cobbs Cavern KFA agreement. Although the 163-acre KFA is not currently fenced, most of the 1,670-acre Cobbs Ranch that surrounds it is fenced. The remote location of the Cobbs Ranch and metal gating on the entrances to Cobbs Cavern ensures that trespassing is not considered an issue. Livestock no longer have access to the KFA. A rock wall near the cave entrance that required modification has been replaced by a French drain and the area has been revegetated with native grasses. A roof extension that was covering the main entrance to the cave has been removed to allow more natural movement of air and rainfall access to the cave entrance. However, the roof removal has allowed increased air circulation at the cave entrance and troglobitic species have moved farther back into the feature recesses to avoid surface climate conditions. Finally, a spill plan and secondary containment unit related to a Pedernales Electric Cooperative (PEC) transformer has not been implemented. Repeated contact with the PEC has resulted in an agreement that the unit will be removed, however the company has not actually done so. Due to the fact that the transformer is not electrified, contains no polychlorinated biphenyls (PCBs), sits at ground level, and is over 1.75 miles from the main entrance to Cobbs Cavern, contamination from the transformer is unlikely.

2.3.2 Hydrogeology

Cobbs Cavern and Cobbs Spring are part of an integrated karst hydrologic system contained within the discrete section of Edwards limestone defined by the fault, which passes roughly 1,800 feet southeast of the spring outlet and erosional removal of the Edwards by tributaries of Salado and Berry Creeks to the northwest of the fault (see Figure 7). The maximum theoretical springshed consists of all rocks that are higher than the stratigraphic position and elevation of the spring due to the unconfined nature of the aquifer. However, in the area surrounding the property, there are numerous incisions into the Edwards outcrop formed by drainages other than Cobbs Springs Branch. It is a predictable regional pattern that significant incision into the Edwards outcrop form groundwater discharge points at the Edwards/Comanche Peak contact, ranging from very low flow seeps supporting slightly more luxuriant vegetation than surrounding areas to permanent springs such as Cobbs Spring. Twelve permanent springs (including several other Georgetown salamander locations) and numerous seeps occur along this lithologic contact within a 10-mile radius of the property. These additional discharge points constrain the boundaries of the actual recharge area that contributes to Cobbs Spring and were subtracted from the idealized springshed shown in Figure 7. A geologic map property was produced from geologic maps compiled at the University of Texas Bureau of Economic Geology and on-site field mapping by SWCA Environmental Consultants (SWCA) (see Figure 7).

Figure 8 presents a cross-sectional illustration of the hydrologic mechanism behind the spring flow. The Comanche Peak is a relatively impermeable unit that perches groundwater working its way down through the overlying Edwards limestone. Groundwater moves laterally along the contact between the two units and downhill (the fault block upon which the recharge area sits is tilting toward the southeast at about 40 feet per mile) until it finds daylight, where the down cutting of Cobbs Springs Branch exposes the Comanche Peak. Spring flow was probably initiated when the bed of Cobbs Springs Branch intersected the Comanche Peak along the fault plane, which is approximately 1,800 feet downstream from the current spring outlet (see Figure 8).




Figure 8. Cobbs Spring cross sectional representation showing hydrologic mechanism behind spring flow.

Headward erosion has caused the spring outlet to gradually move upstream. This is evidenced by the trend of Cobbs Cavern itself. Extending the linear trend of the cavern along its bearing to the northeast produces a line pointing directly to the intersection of the fault plane with the current spring run. The cave, therefore, appears to be a paleo-channel pointing to the location of the paleo-spring outlet. Based on the above understanding of site hydrology, the springshed of Cobbs Spring is approximately 750 acres, with approximately 16% of the springshed occurring within the boundaries of the current conservation easement.

2.3.3 Caves of the Cobbs Cavern KFA

Cobbs Cavern: Cobbs Cavern can be thought of as two caves within one. Cobbs Cavern's eastern third was operated on a limited basis as a show cave in the late 1960s and early 1970s (Photograph 6). The "Show Side" has been made easily traversable with a wide walking path in the cave's center. Visitors have to stoop in a few places, but the cave is otherwise very easy to move through with no tight squeezes or areas that require crawling. Public access to the cave has been restricted since that time, but the old visitor center building remains. The western two-thirds of Cobbs Cavern is called the "Wild Side" and is in a natural state with difficult to travel terrain and muddy, tight squeezes not suitable for novice cave exploration. The Bone Cave harvestman and *Batrises texanus* are known from Cobbs Cavern. A cave gate currently prevents unauthorized access to this feature.

Despite concerns about the potential effects of entrance modification and trail construction associated with tourism activities, the invertebrate fauna of the cave remains quite diverse. Aside from the rare species listed above, the cave community also includes at least two species of *Ceuthophilus* cave crickets, *Cambala* millipedes, *Texoreddellia* silverfish, *Pseudouroctonus* scorpions, *Eidmannella* spiders, *Collembola* (springtails), and mites. Cobbs Cavern is one of the 30 longest caves of more than 6,000 caves in Texas. Cobbs Cavern and associated non-navigable void network is one of the single largest habitats known for endangered karst invertebrates in the State.

Two karst feature excavations west of the mapped end of the cave have revealed troglobitic *Cicurina* spiders and *Speodesmus* millipedes. This indicates that the Cobbs Cavern habitat extends at least

1,000 feet beyond the mapped passage and that additional loci of nutrient input occur in addition to the known entrances at either end of the former tourist trail.

A wildfire burned across most of the conservation easement during the summer of 2008, creating several opportunities for additional investigation. Several significant karst features that were previously hidden by dense brush were located, including an old entrance to the cave and several sinkholes indicating additional passage (and troglobite habitat) beyond the cave's eastern-mapped end during a site inspection following the fire. Shrubby vegetation regrowth following the fire may lead to BCVI habitat creation and potential occupation. Migrating BCVI were sighted on the ranch in 2004.



Photograph 6. Photograph of the interior of Cobbs Cavern.

Cobbs Cavern KFA protection is significant to covered karst invertebrate species recovery and additional species conservation due to likely genetic distinctiveness. Cobbs Cavern is the northern-most documented location for the Bone Cave harvestman, *Batrissodes texanus*, and several additional troglobites (Photographs 7–12). It is also notable as the first known location for the species *Tayshaneta anopica* (Photograph 13). This spider lacks eyes entirely (as the scientific name suggests) and is considered a very advanced troglobite. Two of its close relatives (*Tayshaneta microps* from Bexar County and *T. myopica* from Travis County) are listed as endangered but are not as advanced in their troglomorphy, as they have eyes of reduced size.

In January 2005, SWCA conducted a cursory analysis of the likely drainage area that contributes water to Cobbs Spring to determine how much area would be preserved by conservation bank establishment. SWCA estimated the springshed area using several local geologic parameters. The property's lithological outcrops are Cretaceous in age and consist of the Georgetown limestone, Edwards limestone, and a previously unmapped outcrop of the Comanche Peak limestone (see Figure 7). The Georgetown and Edwards limestones are relatively transmissive units that form an unconfined karstic aquifer perched above the Comanche Peak limestone, which does not support easy movement of groundwater due to

higher marl content. One normal fault crosses the property and produces an abrupt change in surface lithology from the Edwards limestone on the up-thrown block to the Georgetown Limestone on the down-thrown block (see Figures 7 and 8).

Table 6 shows documented species within Cobbs Cavern.



Photograph 7. Bone Cave harvestman collected in Cobbs Cavern.



Photograph 8. *Batrisodes texanus* collected in Cobbs Cavern.



Photograph 9. *Rhadine noctivaga* collected in Cobbs Cavern.



Photograph 10. *Speodesmus bicornourus* collected in Cobbs Cavern.



Photograph 11. *Cicurina vibora* collected in Cobbs Cavern.



Photograph 12. *Eurycea* salamander at Cobbs Spring.



Photograph 13. *Tayshaneta anopica* collected in Cobbs Cavern.

Table 4. Species previously documented from the Cobbs Cavern KFA.

Species		Cobbs Cavern
Permitted Species		
	<i>Texella reyesi</i>	√
	<i>Batrises texanus</i>	√
Other Species		
Crickets	<i>Ceuthophilus</i> sp.	√
Arachnids	<i>Cicurina varians</i>	√
	<i>Cicurina vibora</i>	√
	<i>Tayshaneta anopica</i>	√
	<i>Eidmannella</i> sp.	√
	<i>Achaeearanea</i> sp.	√
	<i>Pseudouroctonus reddelli</i>	√
	<i>Tartarocreagris</i> sp.	√
Millipedes/ Centipedes	<i>Speodesmus bicornourus</i>	√
	<i>Cambala speobia</i>	√
	<i>Oxidus gracilis</i>	√
	<i>Scolopendra</i> sp.	√
Beetles	<i>Rhadine noctivaga</i>	√
Reptiles/ Amphibians	<i>Crotalus atrox</i>	√
Mammals	<i>Bassariscus astutus</i>	√
	<i>Procyon lotor</i>	√
	<i>Perimyotis subflavus</i>	√
	<i>Myotis velifer</i>	√
Other	Isopod	√
	<i>Texoreddellia aquilonalis</i>	√
	Gnats	√
	<i>Collembola</i> sp.	√
	<i>Heliodiscus eigenmanni</i>	√
	Gastropoda	√

2.4 Millennium Preserve

2.4.1 Introduction

Table 5. Millennium Preserve snapshot showing basic preserve information.

Millennium Preserve Snapshot	
Preserve Inception Year	2002
Acreage	74.4
Fence Status	Open-ended fences surround species caves
Sign Status	Signs posted on fences, warning signs on all cave gates
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Through Trip, Little Demon, Millennium, Fence and Trail
Non-gated Caves	Knuckle, Cap, Forest Elms Sink

Williamson County purchased the approximately 800-acre Southwest Williamson County Regional Park (Regional Park) in 2002 with County bond money supplemented by funding from TxDOT. The Regional Park is located northeast of the intersection of Farm-to-Market Road (FM) 1431 and CR 175 near Leander, Texas, and encompasses the Wilco and Millennium Preserves. The Regional Park's developed portion contains recreational facilities including ball fields, tennis courts, a track, disc golf course, and miniature train track. Most Regional Park acreage is open space, including two dedicated preserves for the Bone Cave harvestman and additional species. Other caves occur in open space outside of the karst preserves. The Regional Park occurs within the Georgetown KFR and its two preserves were identified by the USFWS as a potential KFA in the 5-year review of Bone Cave harvestman (USFWS 2009a). The Regional Park occurs adjacent to areas identified in the RHCP as potential habitat for the GCWA, and woodland within the park may become suitable for GCWAs over time. The WCCF headquarters is located within the Regional Park and maintenance staff are able to visit cave entrances on a regular basis.

An undefined portion of the park is recognized by the USFWS as a karst conservation bank. The County received funding from TxDOT in conjunction with the SH 45 Section 7 ESA consultation. These funds contributed to the 129-acre preserve land establishment to protect the ecological integrity of four high-quality caves within the park.

The Regional Park is geologically representative of a broader, highly cavernous area within the northern segment of the EARZ. This cavernous area contains dozens of caves and stretches for approximately 1.5 miles to the south and east. The Regional Park consists primarily of broad, flat uplands drained to the south by Brushy Creek tributaries. Total relief is approximately 60 feet, ranging from 935 to 870 feet amsl.

Mike Warton and Associates (2000) and Horizon Environmental Services (2002) collectively described 39 features including caves, sinkholes, solution cavities, and fracture zones within the Regional Park. All features in the park are formed in the Edwards limestone, which is approximately 80 feet thick in the area. The surface expression of caves and other karst features is found generally, but not exclusively, above 900 feet in elevation, and karst features are rarely expressed within a 1,600-foot-wide zone coincident with the primary drainage channel running northwest to southeast through the undeveloped portion of the Regional Park. Two distinct karst areas are divided by this zone: The Millennium Preserve includes

Millennium, Little Demon and Through Trip Caves (plus four other minor caves); whereas the Wilco Preserve contains Wilco, Wild West, Rock Ridge, and Mongo Caves (plus nine other minor caves or karst features). Figure 9 shows both the Millennium and Wilco Preserves in cross section to demonstrate the geologic separation between them.




Figure 9. Geologic cross section across Millennium and Wilco Preserves highlighting karst erosion by Honey Bear Creek.

The 74.43-acre Millennium Preserve maintains fences containing appropriate signage strategic placed to direct the public away from the many of the caves. However, the fences are open-ended and this is a popular location with many trails designed for public-access hiking. A publicly accessible hike and bike pathway extends through Beck Preserve's western side but does not pass within either the drainage or ecological buffers established for the occupied caves. Cave gates prevent access to Millennium, Little Demon and Through Trip Caves. Figure 10 shows the Millennium Preserve and associated caves.




Figure 10. Millennium Preserve map showing documented caves.

2.4.2 Hydrogeology

Caves within the Millennium Preserve occur on the northeastern flank of a subtle southeast-trending ridge. Millennium Preserve slopes generally to the northeast contributing runoff to the Honey Bear Creek drainage basin, which is a tributary of Brushy Creek. Total relief on Millennium Preserve is approximately 40 feet ranging from 930 to 890 feet amsl. Most of the cave entrances are formed between 920 and 925 feet amsl and most traversable cave passage occurs between 905 and 925 feet amsl (Figure 11). Cave distribution suggests the presence of a well-developed, stratigraphically controlled dissolution zone that likely extends somewhat continuously beneath the ridge. This karstic horizon within the bedrock likely provides mesocavern habitat between the occupied caves and likely beyond.

The Millennium Preserve is entirely underlain by the Edwards limestone, which is the primary cavernous bedrock unit in the region (see Figure 11). The Millennium Preserve is on the EARZ's northern segment and groundwater recharge primarily occurs where the Edwards limestone is exposed at the surface through direct infiltration of precipitation on the limestone outcrop, through streamflow loss, and through secondary porosity features such as faults, fractures, and karst features (caves, solution cavities, sinkholes, etc.). The relatively flat terrain within Millennium Preserve is conducive to slow runoff which enhances the potential for aquifer recharge. Groundwater perches at the base of the Edwards limestone where the underlying Comanche Peak acts as an aquitard and diverts flow laterally.

No faults are known to occur within the Millennium preserve, although fractures are expressed within all caves. A previously mapped fault passes approximately 0.5 mile northwest from the Millennium Preserve (Collins 2005). Field mapping by Cambrian personnel indicates the presence of at least two previously unmapped faults southeast from the Millennium Preserve.

The Edwards limestone is approximately 85 feet thick within the Millennium Preserve (Collins 2005; Senger et al. 1990). The Edwards limestone is near its full thickness near Millennium Preserve's highest topography and is nearly breached along Honey Bear Creek northeast preserve (see Figures 9 and 10). It is this near breach of cavernous strata that provides an isolating mechanism between the Millennium and Wilco Preserves. The Honey Bear Creek basin follows an approximately 1,500-foot-wide erosion corridor karst feature strata is generally absent. Field mapping indicates the Edwards limestone is only 5 feet thick within the basin's southern end and not more than 15 feet thick at the northern end of the corridor. The Edwards limestone pinches out altogether immediately south from the Millennium Preserve and groundwater can be observed discharging from several seeps along Honey Creek at the contact with the underlying Comanche Peak formation. The thin Edwards limestone remnant along the creek bottom area is covered with relatively thick clay soils which are absent from the Millennium Preserve's karstic uplands. Clay soil areas throughout the corridor are marked by mesquite trees and several small stock ponds that hold water after rains for an extended period (depressions on the Edwards typically do not hold water due to the porous nature of the rock). Low permeability, thin outcrop, and the general absence of karst features throughout this corridor indicate that suitable habitat for karst invertebrates is unlikely to occur there.

Surface drainage basin delineation is a relatively straightforward process based on the surface topography near cave entrances and other nearby features such as sinkholes or fractures with the potential to convey water into the cave. All caves within the Millennium Preserve have relatively limited surface drainage areas due to their upland locations. The species-occupied caves all draw their surface runoff from a ridge contained within the Millennium Preserve boundaries. Therefore, all surface runoff reaching the species-occupied cave entrances and footprints originates within the Millennium Preserve.




Figure 11. Millennium Preserve showing geological cross section.

2.4.3 Caves of the Millennium Preserve

Cave maps are included within Appendix C.

Through Trip Cave: The entrance to Through Trip Cave was excavated to 4.5 feet to find a horizontal passage (Photograph 14). A tunnel extended for 6.5 feet before becoming constricted. Further excavation located open passage leading west for approximately 30 feet. A small crawlway was further excavated and continues west for another 20 feet before leading to another small crawlway. This crawlway was excavated and led to another room continuing west. This opens immediately into a 7-foot-high passage and continues west for 40 feet before leading to another shaft within the rock joint. This shaft was revealed to be a second cave entrance. The surveyed cave length is approximately 105.9 feet, with a 16-foot maximum depth. The cave footprint measures approximately 80 feet long by 32 feet wide. This cave is not currently known to contain any endangered karst invertebrates and a cave gate prevents unauthorized access.



Photograph 14. Through Trip Cave west entrance.

Little Demon Cave: The Little Demon Cave entrance was excavated to 6 feet, where an additional 6-foot open drop was found (Photograph 15). Open cave passage was encountered at 12 feet. The mostly rock joint-controlled room trends N90W and contains several passages in various directions. The surveyed cave length measures approximately 75.6 feet, with a 19-foot maximum depth. The cave footprint measures approximately 45 feet long by 14 feet wide. This cave is known to contain the Bone Cave harvestman. A cave gate currently prevents unauthorized access to this feature.



Photograph 15. Little Demon Cave entrance.

Millennium Cave: This cave had the only open entrance within the Regional Park that required no digging (Photograph 16). The entrance room is approximately 28 feet long, and at the end, a digging lead was pursued. A large room was encountered at 6 feet deep after additional excavation. This room measures approximately 40 feet in diameter with a floor-to-ceiling height of 4.5 feet. The surveyed cave length is approximately 82.7 feet, with a 29-foot maximum depth. The cave footprint measures approximately 80 feet long by 35 feet wide. This cave is known to contain the Bone Cave harvestman. A cave gate currently prevents unauthorized access to this feature.



Photograph 16. Entrance to Millennium Cave.

The Millennium Preserve contains four additional caves and sinkholes not currently known to contain habitat for troglobitic fauna. This is primarily due to their small size and inaccessibility for researchers. However, the collective karst feature distribution suggests the Millennium Preserve's cave containing strata are an interconnected cavernous and mesocavernous habitat area spanning more than 20 acres. The cave geomorphology suggests they formed from a larger underlying karst void network collapse. That network likely formed under phreatic conditions in response to paleo-aquifer karst hydrology as groundwater followed an easterly to northeasterly path toward the confined aquifer zone. This process was initiated in the Miocene following the tectonic events that created the Balcones Escarpment (Senger et al. 1990; White 2009). Aquifer entrenchment and regional streams caused the water table to drop below the current cave elevation. The buoyant force acting on the cave ceilings was lost and the caves began to collapse when this happened. The cave floors that are accessible now are largely formed from breakdown material that has collapsed from the original cave ceiling. The modern cave network has chaotic morphology relative to the paleo cave network because the collapse process is incomplete. What formed as a relatively continuous void network formed in response to dissolution by flowing groundwater is now a seemingly discontinuous patchwork of air pockets from a human cave explorer's perspective. However, the mesocavern network is far more extensive and accessible from a karst invertebrate's perspective, which still emulate the original groundwater pathways. These species are referred to as karst invertebrates rather than cave invertebrates due to their occupancy of the fabric of the aquifer and not just the larger cavernous spaces.

Fence and Trail Cave: The entrance to Fence and Trail Cave was excavated to 7 feet deep where open passage was discovered (Photograph 17). The passage is formed along a rock joint trending N70E. The entrance extends through a tight constriction into a passage approximately 5 feet wide by 13 feet long, with 2-foot maximum floor to ceiling height. The surveyed cave length measures approximately 20 feet, with a maximum depth of 7 feet. The cave footprint measures approximately 20 feet by 5 feet. This cave

is not currently known to contain any endangered karst invertebrates, is not monitored for karst biota, and has a cave gate to prevent unauthorized access.



Photograph 17. Entrance to Fence and Trail Cave.

Knuckle Cave: The Knuckle Cave entrance was excavated to 5 feet, where a small open cave was encountered (Photograph 18). The feature appears to be formed along a rock joint trending N30W. The cave extends to the southeast along a low bedding plane for about 20 feet. The surveyed cave length is approximately 29.2 feet, with a maximum depth of 7.3 feet. The cave footprint measures approximately 26 feet long by 11 feet wide. This cave is not currently known to contain any endangered karst invertebrates, is not surveyed for biota, and does not have a gate.



Photograph 18. Entrance to Knuckle Cave.

Cap Cave: Cap Cave consists of a large single solution cavity discovered during the Regional Park development. The entire footprint of the irregularly shaped feature is less than 15 feet in diameter and approximately 5 feet deep. The name “Cap” cave comes from the large flat boulder currently covering the entrance of the feature. No Cap Cave map is available. This cave is not currently known to contain any endangered karst invertebrates, is not surveyed for biota, and does not have a gate.

Forest Elms Sink: Forest Elms Sink is so named due to the presence of a small grove of cedar elm trees, the largest of which have died since the feature was excavated in 2000. The feature consists of a small pit measuring approximately 5 feet in diameter. It was originally mapped as being up to 8 feet deep, but it appears that some slumping from the walls has filled the lower portion of the feature since it was excavated. This cave is not currently known to contain any endangered karst invertebrates, is not surveyed for biota, and does not have a gate.

Table 8 shows species documented from the Millennium Preserve.

Table 6. Species previously documented from the Millennium Preserve.

Species		Wild West Cave	Mongo Cave	Rock Ridge Cave	Wilco Cave	Prospector	Venture
Permitted Species							
<i>Texella reyesi</i>		√	√	√	√		
Other Species							
Crickets	<i>Ceuthophilus secretus</i>	√	√	√		√	√
	<i>Ceuthophilus</i> sp.	√	√	√	√	√	√
Arachnids	<i>Cicurina vibora</i>	√	√	√		√	√
	<i>Cicurina varians</i>	√	√	√	√	√	√
	<i>Anapistula secreta</i>		√				
	Araneae (General Spider)	√		√			
	<i>Achaearanea</i> sp.		√			√	√
	<i>Leiobunum</i> sp.				√	√	√
	<i>Tartarocreagris infernalis</i>		√				
Millipedes/ Centipedes	<i>Speodesmus bicornourus</i>		√		√		
	<i>Cambala speobia</i>	√	√	√	√		
	<i>Scutigera</i> sp.		√			√	√
	<i>Scolopendra</i> sp.						√
Beetles	<i>Rhadine subterranea</i>		√				
	Staphylinidae					√	
	<i>Batrissodes unicomis</i>		√				√
Reptiles/ Amphibians	<i>Eleutherodactylus marnockii</i>		√		√		√
	<i>Crotalus atrox</i>		√				
	<i>Lithobates berlandieri</i>		√				
Mammals	<i>Perimyotis subflavus</i>			√		√	√
	<i>Myotis velifer</i>		√				
Other	Desert Cockroach	√	√	√			
	Isopod	√	√	√	√		
	Assassin Bugs	√	√				√
	<i>Heliodiscus eigenmanni</i>		√	√		√	√
	Bark Louse	√					
	Earthworm		√			√	
	Gnat					√	
	<i>Collembola</i> sp.	√	√	√	√	√	√

2.5 Wilco Preserve

2.5.1 Introduction

Table 7. Wilco Preserve snapshot showing basic preserve information.

Wilco Preserve Snapshot	
Preserve Inception Year	2002
Acreage	152.5
Fence Status	Open-ended fences surround endangered species caves
Sign Status	Signs posted on fences, warning signs on all cave gates
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Wilco, Rock Ridge, Mongo, Wild West, Choya, Talus, No Paseo, Nuevo, Prospector, Venture
Non-gated Caves	Poppin Rocks, Circulation Sink, Lockout Sink, Side Pocket Sink, West Boundary Sink

The Wilco Karst Preserve is located within the Regional Park near Leander, Texas (Figure 12) and was acquired by Williamson County under the same pretense as the Millennium Preserve. The flora and fauna have begun a successional transition from overgrazed, managed grassland to a predominantly native woodland/grassland mosaic from ranching cessation more than 15 years ago and provides suitable habitat for native species. The Wilco Preserve occurs adjacent to areas identified in the RHCP as potential habitat for the GCWA and woodland within the park may become suitable for GCWAs over time (SWCA et al. 2008).




Figure 12. Wilco Preserve map showing documented karst features.

2.5.2 Hydrogeology

Total relief on the Wilco Preserve is approximately 60 feet ranging from approximately 935 to 875 feet amsl. Land slopes generally from uplands in the northeast to drainages exiting the western and southern Wilco Preserve boundaries. The Honey Bear Creek drainage basin is immediately west from the Wilco Preserve and separates it from the Millennium Preserve. The Wilco Preserve is located entirely within the EARZ. Based on regional patterns, recharge from the property likely follows the groundwater gradient to the northeast, but no formal study has been conducted.

The Wilco Preserve caves with documented Bone Cave harvestman presence occur on either flank of a subtle southwest-trending ridge. Most cave entrances are formed between 910 and 925 feet amsl and most traversable cave passage occurs between 905 and 890 feet amsl (see Figure 9). This distribution suggests the presence of a highly developed, stratigraphically controlled dissolution zone which likely extends somewhat continuously beneath the ridge. This karstic horizon within the bedrock likely provides mesocavern habitat between the four occupied caves and likely beyond. The Edwards limestone is approximately 85 feet thick around Wilco Preserve (Collins 2005; Senger et al. 1990). Accordingly, the Edwards is near its full thickness near the highest topography and is nearly breached along Honey Bear Creek south from Wilco Preserve (see Figure 9). This near cavernous strata breach provides an isolating mechanism between the Wilco and Millennium Preserves. No faults are known to occur within Wilco Preserve although fractures related to regional faulting are expressed within all of the caves. A mapped fault passes approximately 0.5 mile northwest of the Wilco Preserve (Collins 2005).

The Wilco Preserve caves have relatively limited surface drainage areas because of their upland terrain location. The species-occupied caves all draw their surface runoff from the ridge described above. Therefore all surface runoff reaching the cave entrances and footprints originates from within the Wilco Preserve. The relatively flat terrain within the Wilco Preserve is conducive to slow runoff, which enhances the potential for aquifer recharge.

2.5.3 Caves of the Wilco Preserve

Cave maps are included within Appendix C.

The Wilco Karst Preserve contains Wilco Cave, Wild West Cave, Mongo Cave, and Rock Ridge Cave (all of which contain the Bone Cave harvestman), plus 10 additional significant karst features. Wilco Preserve contains the endangered Bone Cave harvestman along with a representative sampling of local endemic cave fauna including the troglobitic species *Cicurina browni*, *Tartarocreagris infernalis*, *Speodesmus bicornourus*, and *Rhadine subterranea mitchelli*, which are additional species addressed in the RHCP (SWCA et al. 2008). Additional fauna are likely to be detected in each cave now that regular biological monitoring is being conducted (Krejca and Weckerly 2007; Schneider and Culver 2004). Most Wilco Preserve caves are likely quite small relative to the volume of potential mesocavern habitat around and between the caves. A dominant east/west fracture trend is expressed in the morphology of all of the caves.

Wild West Cave: The Wild West Cave entrance was excavated to 4 feet and an open passage was encountered. The entrance opens to the west into a long linear room measuring approximately 18 feet wide by 40 feet long, with a 5.5-foot maximum floor to ceiling height. Access to a second lower-level room continues west from the main room's northwest corner. The lower-level room is wide and low and measures approximately 25 feet long by 35 feet wide, with a 3-foot maximum floor to ceiling height. The surveyed cave length is approximately 88.4 feet, with a 15.5-foot maximum depth. The cave footprint measures approximately 60 feet long by 38 feet wide. The cave is formed along a rock joint trending

N90W and is known to contain the Bone Cave harvestman. A cave gate currently prevents unauthorized access to this feature.

Mongo Cave: Mongo Cave is located within the tract's largest sinkhole, which measures approximately 75 feet long by 50 feet wide by 6 feet high. The sinkhole is formed along a rock joint trending N90W and funnels down to a main drainage portal that was excavated to 3 feet before encountering open cave passage. The excavated entrance portal descends at a sharp angle for about 20 feet into a large single room measuring approximately 86 feet long by 41 feet wide, with a 6-foot maximum floor to ceiling height. A 12-foot-deep pit was excavated in this main room and a small lower-level room was located. No further extent was found. The surveyed cave measures approximately 167.8 feet long, with a maximum depth of 29.3 feet. The cave footprint measures approximately 86 feet long by 41 feet wide. Mongo Cave is the single largest cave known from the tract. This cave is known to contain the Bone Cave harvestman. A cave gate currently prevents unauthorized access to this feature.

Rock Ridge Cave: The Rock Ridge Cave entrance was excavated to 4.5 feet and open passage was located. The cave entrance and extent occur along a rock joint trending N85E. The cave extends primarily west for approximately 60 feet until passage becomes blocked. The cave also extends east from the entrance for approximately 20 feet. The eastern passage extends beneath the property boundary fence of the adjacent property. The surveyed cave length measures approximately 103.7 feet long, with a 14-foot maximum depth. The cave footprint measures approximately 80 feet long by 14 feet wide. Rock Ridge Cave is known to contain the Bone Cave harvestman. A cave gate currently prevents unauthorized access to this feature.

Wilco Cave: Wilco Cave (Photograph 19) was excavated to 9 feet and semi-open passage was encountered. A short passage leads west and drops down to a small room. A low bedding plane void extends south for an undetermined distance within this small room. The surveyed cave length measures approximately 47 feet long, with a 15.7-foot maximum depth. The cave footprint measures approximately 37 feet long by 27 feet wide. Wilco Cave is known to contain the Bone Cave harvestman. A cave gate currently prevents unauthorized access to this feature.



Photograph 19. Entrance to Wilco Cave.

Prospector Cave: Prospector Cave is formed along a rock joint trending east 50 feet, with a 19-foot maximum depth. Further passage occurs below a drain and beyond restriction in the cave's terminal east end. Prospector Cave is formed within the same karstic horizon as the species-occupied caves and may represent an exposure of the occupied mesocavern habitat. It is especially close to Mongo Cave and may be directly connected. This cave is not currently known to contain any endangered karst invertebrates, is currently monitored for karst biota, and has a cave gate to prevent unauthorized access to this feature.

Venture Cave: Venture Cave is formed along a rock joint trending northeast for approximately 70 feet, with a 21-foot maximum depth. Further passage occurs below a drain and beyond an air blowing restriction in the cave's terminal northeast extent. This cave is not currently known to contain any endangered karst invertebrates, is currently monitored for karst biota, and has a cave gate to prevent unauthorized access to this feature.

Talus Cave: Talus Cave is within an approximately 40-foot diameter sinkhole that funnels down to a single portal. The portal was excavated to 9 feet where open passage was located. A constricted passage continued down at a sharp angle to 24 feet and a humanly accessible, cone-shaped collapse extends for approximately 15 feet. The surveyed cave length cave measures approximately 67.3 feet long, with a 24-foot maximum depth. The cave footprint measures approximately 43 feet long by 20 feet wide. This cave is not currently known to contain any endangered karst invertebrates, is not monitored for karst biota, and has a cave gate to prevent unauthorized access.

No Paseo Cave: No Paseo cave was excavated 5 feet deep to reveal open passage extending both southwest and northeast. The passage extends approximately 20 feet from the cave entrance northeast and ends at a low bedding plane. The passage extends approximately 5 feet southeast before declining to an approximately 7-inch diameter drain that extends down vertically for about 7 feet. The surveyed cave length measures approximately 36.5 feet, with a 13.5-foot maximum depth. The cave footprint measures

approximately 26 feet long by 11 feet wide. This cave is not currently known to contain any endangered karst invertebrates, is not monitored for karst biota, and a cave gate currently prevents unauthorized access.

Choya Cave: Choya Cave was excavated to 5 feet where open passage formed along a rock joint trending N40E (Photograph 20). The passage extends northeast for approximately 35 feet. The passage then extends northwest to an approximately 30-foot-long by 1-foot-wide room, with 4.5-foot maximum floor to ceiling height. The surveyed cave length measures approximately 155.8 feet long, with an 8.4-foot maximum depth. The cave footprint measures approximately 41 feet long by 31 feet wide. This cave is not currently known to contain any endangered karst invertebrates, is not monitored for karst biota, and has a cave gate to prevent unauthorized access.



Photograph 20. Entrance to Choya Cave.

Poppin Rocks Cave: Poppin Rocks Cave was excavated 10 feet, where a larger cave structure was revealed. However, the passage is blocked by massive infill and collapse material. The accessible cave extent measures approximately 18.5 feet long, with a 10-foot maximum depth. This cave is not currently known to contain any endangered karst invertebrates, is not surveyed for biota and does not have a gate.

Nuevo Cave: Nuevo Cave was excavated to depth 4 feet, where a small, very low bedding plane passage was encountered. Cave extent follows a rock joint trending N90W into a very low room just a few feet below the ground surface level. The surveyed cave length measures approximately 21.3 feet long, with a 539-foot maximum depth. The cave footprint measures approximately 28 feet long by 10 feet wide. This cave is not currently known to contain any endangered karst invertebrates, is not monitored for karst biota, and has a cave gate to prevent unauthorized access to this feature.

Circulation Sink: This sinkhole was enlarged sufficiently (Photograph 21) to allow entry to a small chamber with a low bedding plane void extending northwest for approximately 15 feet. Airflow comes

from an additional cave passage that is not currently accessible. This feature is not known to contain endangered karst invertebrates, is not currently monitored for karst biota, and does not have a cave gate.



Photograph 21. Entrance to Circulation Sink.

Lockout Sink: This sinkhole was excavated 5 feet to expose a 3-inch-diameter solution cavity with airflow issuing from a low bedding plane. This feature is not known to contain endangered karst invertebrates, is not currently monitored for karst biota, and does not have a cave gate.

Side Pocket Sink: This sinkhole was enlarged sufficiently to allow entry to a small chamber measuring approximately 5 feet in diameter by 3 feet high with a low bedding plane void extending north for approximately 15 feet. Side Pocket Sink is formed within the same karstic horizon as the species-occupied caves and may represent occupied mesocavern habitat exposure, which could be an important nutrient input point between Wilco and Rock Ridge Caves. This feature is not monitored for karst invertebrates and does not have a cave gate.

West Boundary Sink: This sinkhole was excavated 4 feet to expose a bedding plane void extending to the east with airflow. The feature is very close to Wild West Cave and may represent mesocavern habitat connected to Wild West Cave. West Boundary Sink is formed within the same karstic horizon as the species-occupied caves and may represent occupied mesocavern habitat exposure. This feature is not monitored for karst invertebrates and does not have a cave gate.

Table 10 shows documented species within the Wilco Preserve.

Table 8. Species previously documented from the Wilco Preserve.

Species		Wild West Cave	Mongo Cave	Rock Ridge Cave	Wilco Cave	Prospector	Venture
Permitted Species							
<i>Texella reyesi</i>		√	√	√	√		
Other Species							
Crickets	<i>Ceuthophilus secretus</i>	√	√	√		√	√
	<i>Ceuthophilus</i> sp.	√	√	√	√	√	√
Arachnids	<i>Cicurina vibora</i>	√	√	√		√	√
	<i>Cicurina varians</i>	√	√	√	√	√	√
	<i>Anapistula secreta</i>		√				
	<i>Tartarocreagris infernalis</i>		√				
	Araneae	√		√			
	<i>Achaeearanea</i> sp.		√			√	√
	<i>Leiobunum</i> sp.				√	√	√
Millipedes/ Centipedes	<i>Speodesmus bicornourus</i>		√		√		
	<i>Cambala speobia</i>	√	√	√	√		
	<i>Scolopendra</i> sp.						√
	<i>Scutigera</i> sp.		√			√	√
Beetles	Staphylinidae					√	
	<i>Batrissodes uncinornis</i>		√				√
	<i>Rhadine subterranea</i>		√				
Reptiles/ Amphibians	<i>Crotalus atrox</i>		√				
	<i>Lithobates berlandieri</i>		√				
	<i>Eleutherodactylus marnockii</i>		√		√		√
Mammals	<i>Perimyotis subflavus</i>			√		√	√
	<i>Myotis velifer</i>		√				
Other	Desert Cockroach	√	√	√			
	Isopod	√	√	√	√		
	Assassin Bugs	√	√				√
	<i>Heliodiscus eigenmanni</i>		√	√		√	√
	<i>Collembola</i> sp.	√	√	√	√	√	√
	Bark Louse	√					
	Earthworm		√			√	
	Gnat					√	

2.6 Chaos Cave Preserve

2.6.1 Introduction

Table 9. Chaos Cave Preserve snapshot showing basic preserve information.

Chaos Preserve Snapshot	
Preserve Inception Year	2008
Acreage	30.0
Fence Status	Fully fenced
Sign Status	Signs posted around perimeter
Baseline Vegetation Survey Date	None
Owner	Easement- TxDOT
Gated Caves	None
Non-gated Caves	Under-the-Fence, Chaos, Poison Ivy, Rather Gaping Pit

The approximately 30-acre Chaos Preserve is immediately south from SH 45 (Figure 13). It was established by TxDOT as a conservation measure related to potential Bone Cave harvestman impacts from SH 45 construction and Williamson County assumed management responsibility for Chaos Preserve in 2008. The County has agreed to preserve and protect the natural, scenic, open space, and ecological features in a contract for mitigation services between Williamson County and TxDOT dated 26 August 2008. Williamson County will also provide funding for management and monitoring activities and will report on those activities to TxDOT biannually.

Chaos Preserve is bounded by SH 45 to the north, by a rail line to the southwest, and by undeveloped portions of the Robinson Ranch to the east and west (see Figure 13). This area is a relatively flat upland drained to the north by Lake Creek. Elevations range from approximately 800 to 815 feet amsl (Figure 14).

Chaos, Under the Fence, and Poison Ivy Caves are confirmed as containing the Bone Cave harvestman. Karst invertebrates documented within the Chaos Preserve are included in Table 12. Immediately east from the Chaos Preserve are several additional caves, including Sam Bass Hideaway Cave, a known site for the Bone Cave harvestman. Rather-Gaping-Pit is a small feature with little known biological significance and is not monitored for karst invertebrates. The Chaos Preserve was identified by the USFWS as a potential KFA in the Bone Cave harvestman 5-year review (USFWS 2009a). Chaos Preserve is fenced and has appropriate signage to aid trespassing prevention per guidelines set in the RHCP by USFWS.

2.6.2 Hydrogeology

Chaos Preserve is located entirely within the EARZ. Based on regional patterns, recharge from the property likely follows the groundwater gradient to the northeast and east toward Rattan Creek and Smith Lake, but no formal study has been conducted.



Figure 13. Chaos Preserve location map.




Figure 14. Geologic cross section of the Chaos Preserve.

2.6.3 Caves of the Chaos Preserve

Access to the Chaos Preserve is relatively difficult; therefore, no caves are currently gated. Cave maps are included in Appendix D.

Chaos Cave: The Chaos Cave entrance is located within a sinkhole approximately 9 feet in diameter (Photograph 22). The entrance drops vertically for 5 feet and opens into a single room passage measuring approximately 18 feet in diameter. The open cave passage maximum depth is approximately 9 feet. This cave is not currently gated and is known to contain the Bone Cave harvestman.



Photograph 22. Chaos Cave entrance.

Under the Fence Cave: The Under the Fence Cave entrance is located directly beneath the fence separating the Chaos Preserve from the adjacent property (Photograph 23). The cave entrance drops vertically 5 feet and slopes downwards and south. The passage extends approximately 15 feet and terminates in a single room approximately 18 feet long by 9 feet wide. The open cave passage maximum depth is approximately 10 feet. This cave is known to contain the Bone Cave harvestman and does not have a cave gate.



Photograph 23. Under the Fence Cave entrance.

Poison Ivy Cave: The Poison Ivy Cave entrance measures approximately 5 feet long by 1.5 feet wide and slopes down vertically 6 feet (Photograph 24). Mapped cave extent is limited to a single room measuring approximately 18 feet long by 9 feet wide. This cave is known to contain Bone Cave harvestman and does not have a cave gate.



Photograph 24. Poison Ivy Cave entrance.

Rather Gaping Pit: No specific description or map was available for this feature. This feature is not monitored for karst biota, nor does it have a cave gate to prevent unauthorized access.

Table 12 shows documented species within the Chaos Cave Preserve.

Table 10. Species previously documented from the Chaos Cave Preserve.

Species		Chaos Cave	Under the Fence Cave	Poison Ivy Cave
Permitted Species				
<i>Texella reyesi</i>		√	√	√
Other Species				
Crickets	<i>Ceuthophilus cunicularis</i>	√		√
	<i>Ceuthophilus</i> sp.	√	√	√
Arachnids	<i>Cicurina buwata</i>			√
	<i>Cicurina varians</i>	√	√	√
	<i>Tayshaneta</i> sp.	√		
	<i>Eperigone albula</i>	√		
	<i>Agyneta llanoensis</i>	√		
	<i>Eidmannella</i> sp.	√		
	Araneae	√	√	√
	<i>Achaearanea</i> sp.	√	√	√

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan

Species		Chaos Cave	Under the Fence Cave	Poison Ivy Cave
	<i>Vonones</i> sp.			√
	<i>Leiobunum</i> sp.		√	√
	Tick	√		
Millipedes/ Centipedes	<i>Cambala speobia</i>	√		
	<i>Speodesmus bicornourus</i>	√		
	<i>Scutigera</i> sp.	√		√
	<i>Scolopendra</i> sp.	√		√
	<i>Oxidus gracilis</i>			√
Beetles	Coleoptera		√	
	<i>Batrissodes unicoloris</i>	√		
	Staphylinidae	√		√
	<i>Rhadine subterranea</i>	√		
Reptiles/ Amphibians	<i>Eleutherodactylus marnockii</i>	√		√
	<i>Plethodon albagula</i>	√		
	<i>Lithobates berlandieri</i>	√		
Mammals	<i>Perimyotis subflavus</i>	√		
	<i>Myotis velifer</i>	√		
Other	<i>Coragyps atratus</i>	√		√
	Gnats			√
	<i>Collembola</i> sp.	√	√	√
	Black Ant			√
	Earthworm	√		√
	<i>Heliodiscus eigenmanni</i>	√		
	Gastropod			√
	Desert Cockroach			√
	Assassin Bug			√
	Diptera	√	√	√
	Isopod	√	√	√

2.7 Big Oak Cave Preserve

2.7.1 Introduction

Table 11. Big Oak Cave Preserve snapshot showing basic preserve information.

Big Oak Cave Preserve Snapshot	
Preserve Inception Year	2008
Acreage	10.0
Fence Status	None
Sign Status	Warning signs on all cave gates
Baseline Vegetation Survey Date	None
Owner	Easement- TxDOT
Gated Caves	Big Oak Cave
Non-gated Caves	None

Williamson County assumed management responsibility for the Big Oak Cave Preserve in 2008 from TxDOT's Texas Turnpike Authority Division. The Big Oak Preserve is limited to known Big Oak Cave extent and is the only cave currently managed by the County known to contain the Tooth Cave ground beetle (*Rhadine persephone*). TxDOT had originally agreed to Big Oak Cave biological monitoring as a conservation measure related to the 2001 U.S. Highway (U.S.) 183A project's Environmental Impact Statement. The cave occurs within a roughly 0.5-mile-long median area between the "new" U.S. 183A and the "old" U.S. 183 facilities (also known as South Bell Boulevard); with a 225-foot maximum width within a 10-acre area (Figure 15). The cave's footprint and surface drainage area have been avoided by construction, but impervious cover exists above some subsurface drainage basin, which has reduced the cave's capacity to contribute to species recovery. TxDOT funded annual Big Oak Cave for the first four years (2004–2007) following monitoring plan approval in order to study the long-term roadway development and operation on cave species impacts.

The County has agreed to preserve and protect the natural, scenic, open space, and ecological features of Big Oak Cave, to provide funding for monitoring activities, and to report on those activities to TxDOT biannually in a contract for mitigation services between Williamson County and TxDOT dated 26 August 2008. The Big Oak Cave Preserve is not fenced but does maintain two cave gates to prevent unauthorized access to the both cave entrances.

2.7.2 Hydrogeology

Big Oak Cave Preserve is located within the EARZ. Based on the regional hydrologic gradient, cave recharge likely discharges to the northeast along Brushy Creek, but no formal study has been conducted.




Figure 15. Big Oak Cave preserve location map.

2.7.3 Caves of the Big Oak Cave Preserve

Big Oak Cave: No specific size and configuration information for this cave is available. However, Big Oak Cave has two gated entrances (east and west) leading to small, shallow single rooms that are not connected by humanly accessible passages. Photograph 25 shows the western gated entrance. A cave gate currently prevents unauthorized access to this feature.

Table 14 shows documented species within the Big Oak Cave Preserve.



Photograph 25. Big Oak Cave western entrance.

Table 12. Species previously documented from the Big Oak Cave Preserve.

Species		Big Oak Cave
Permitted Species		
<i>Rhadine persephone</i>		√
Other Species		
Crickets	<i>Ceuthophilus</i> sp.	√
Spiders	<i>Cicurina varians</i>	√
	Araneae	√
	<i>Achaeearanea</i> sp.	√
	<i>Cambala speobia</i>	√
Millipedes/ Centipedes	<i>Speodesmus bicornorus</i>	√
	Staphylinidae	√
Reptiles/ Amphibians	<i>Eleutherodactylus marnockii</i>	√
Other	<i>Coragyps atratus</i>	√
	Moth	√
	<i>Collembola</i> sp.	√
	<i>Heliodiscus eigenmanni</i>	√
	Assassin Bug	√
	Isopod	√

2.8 Priscilla's Well Karst Fauna Area

2.8.1 Introduction

Table 13. Priscilla's Well KFA snapshot showing basic preserve information.

Priscilla's Well KFA Snapshot	
Preserve Inception Year	2008
Acreage	51.5
Fence Status	Fully fenced
Sign Status	Warning signs on all cave gates
Baseline Vegetation Survey Date	2007
Owner	Williamson County
Gated Caves	Priscilla's Well, Priscilla's
Non-gated Caves	None

The Priscilla's Well Preserve was the first KFA recognized by the USFWS as contributing to a listed karst invertebrate species recovery. The Priscilla's Well KFA is 51.5 acres and is generally located between Ronald W. Reagan Boulevard and a residential portion of the Sun City development (Figure 16). Priscilla's Well Cave is documented to contain both the Bone Cave harvestman and *Batrisesodes cryptotexanus*; whereas Priscilla's Cave is documented to have the Bone Cave harvestman (Verdoorn 1994). The KFA entirely encompasses a 13.4-acre area proposed as a preserve in the 1994 management plan for Sun City Georgetown (Verdoorn 1994). Priscilla's Well Cave was biologically monitored for five years between 1995 and 2000, and *Batrisesodes cryptotexanus* was found on nine out of 16 sampling occasions. Priscilla's Cave was not biologically monitored beyond the initial Sun City investigation. *Batrisesodes cryptotexanus* may also be found within Priscilla's Cave with continued monitoring given the two cave's proximity and similarity. The Priscilla's Well KFA is generally in "natural" condition, consisting of a mixed woodland/grassland mosaic. Cattle grazing at this location has officially ceased; however, few neighboring animals do make it onto the Priscilla's Well KFA through fence breaks. These incidents are relatively rare and are rectified as soon as possible. Fencing and signage are placed to restrict cave and KFA access. Cave gates are present on both Priscilla's Well Cave and Priscilla's Cave.

Priscilla's Well KFA proximity to Ronald Reagan Boulevard has prompted concern that a spill or other catastrophic event adjacent to the preserve could negatively impact the cave ecosystem contained within the property. However, the road itself is down gradient from the both caves and guardrails have been installed to prevent vehicles carrying fuel (or other contaminants) from accidentally crashing into Priscilla's Well KFA; therefore, impact from such an event is unlikely. Photograph 26 shows three existing best management practices along Ronald Reagan Boulevard adjacent Priscilla's Well KFA: KFA fence (right), rock gabions to slow stormwater moving off Ronald Reagan Boulevard (center), and guardrails to help prevent vehicular accidents from negatively impacting the Priscilla's Well KFA (upper left).




Figure 16. Priscilla's Well KFA location map.



Photograph 26. Three visible best management practices at the Priscilla's Well KFA.

2.8.2 Hydrogeology

Priscilla's Well KFA encompasses a broad, flat hilltop whose highest elevation is slightly more than 940 feet amsl. Both Priscilla's Cave and Priscilla's Well Cave entrances lie within the 940-foot contour line. Neither cave, nor any karst features within the KFA, likely drain a significant area due to the lack of relief from hilltop placement. The features generally drain a few hundred square feet up-gradient from cave entrances.

The Priscilla's Well KFA is underlain by the Edwards Formation which is estimated to range from 120 to 200 feet in regional thickness (Collins 2002; Verdoorn 1994; Senger et al. 1990). There are identifiable preferential development horizons for karst features within the Edwards Formation that correspond with less resistant limestone beds even though the Edwards Formation has not been stratigraphically subdivided north of Austin. Hundreds of karst feature analyses on more than 5,000 acres surrounding the Priscilla's Well KFA (Verdoorn 1994) found that cave development was largely confined to three distinct solution zones at the top, middle, and base of the Edwards respectively. The hilltop containing the Priscilla's Well KFA is capped by an outcrop of the upper solution zone that is isolated by erosion along the drainage divide between Cowan and Berry creeks. All known karst features within the KFA are confined within that zone, which appears to be roughly 20 feet thick.

The Edwards Formation also hosts nearly impermeable limestone and chert alternating beds that serve as barriers to downward moving groundwater creating wet weather seeps where lateral groundwater movement is expressed as hillside seeps which only flow after heavy rain. Alternating zones of dissolution and impermeable beds constrain the most mapped caves vertical extent in the area. Vertical shaft passages are nearly absent in Williamson County except in caves adjacent to faults. No faults are known to occur within the Priscilla's Well KFA. All Priscilla's Well KFA caves are formed

predominantly along bedding planes horizontally with relatively minor vertical expression at their entrances. Surface water entering the caves and karst features within the Priscilla's Well KFA is incorporated with a perched groundwater system where short residence time water drained by gravity radially away from the hilltop to discharge along the hillside. The groundwater flows downward through the upper Edwards Formation solution zone containing the macroscopic karst features until it intersects an impermeable layer that slow progress until exiting the hillside. This interpretation is strongly supported by several wet weather seeps presence at the 887-foot contour line. These seeps were observed flowing following unusually rainy weather in summer 2007 and were not observed during dry periods in 2008.

2.8.3 Caves of the Priscilla's Well KFA

Priscilla's and Priscilla's Well Caves are small voids reflecting the broader fabric and mesocavernous nature of the local karst. The true measure of troglobite habitat within the Priscilla's Well KFA is more accurately represented by karst feature distribution. Cave maps are included within Appendix E.

Priscilla's Well Cave: Priscilla's Well Cave entrance drops 4.5 feet to a ledge-like landing that drops again to 15 feet deep. The cave interior is 19.9 feet high at its highest point and has an irregular but generally oblong shape. The cave measures 68.5 feet long. Biological surveys conducted between 1995 and 2000 found *Batrachoseps cryptotexanus*. Further investigation will likely result in other troglitic species detection. A cave gate currently prevents unauthorized access to this feature.

Priscilla's Cave: Priscilla's Cave has two narrow, vertical entrances that drop down into the main cavern. The larger entrance drops 10.5 feet and the second entrance drops 8.7 feet. The interior is a low, wide circular room, measuring roughly 15 to 20 feet in diameter. Priscilla's Cave was not biologically monitored beyond the initial Sun City investigation and—given the proximity and similarity of the two caves—the mold beetle may be found with continued monitoring. An SWCA biospeleologist documented the Bone Cave harvestman on 8 September 2007 within Priscilla's Cave. A cave gate currently prevents unauthorized access to this feature.

Table 16 shows documented species within the Priscilla's Well KFA.

Table 14. Species previously documented from the Priscilla's Well KFA.

Species		Priscilla's Cave	Priscilla's Well Cave
Permitted Species			
<i>Texella reyesi</i>		√	√
<i>Batrises cryptotexanus</i>			√
Other Species			
Crickets	<i>Ceuthophilus secretus</i>	√	√
	<i>Ceuthophilus</i> sp.	√	√
Arachnids	<i>Cicurina varians</i>	√	√
	<i>Cicurina vibora</i>	√	√
	<i>Tayshaneta</i> sp.		√
	<i>Achaeearanea</i> sp.	√	√
	Araneae	√	√
	<i>Leiobunum</i> sp.		√
	<i>Pseudouroctonus reddelli</i>	√	
Millipedes/ Centipedes	<i>Scutigera</i> sp.		√
	<i>Scolopendra</i> sp.	√	
Beetles	Staphylinidae		√
Reptiles/ Amphibians	<i>Eleutherodactylus marnockii</i>	√	
	<i>Incilius valliceps</i>		√
Mammals	<i>Myotis velifer</i>	√	
Other	<i>Collembola</i> sp.	√	√
	Desert Cockroach		√
	Isopod	√	√
	Mosquito		√

2.9 Woodland Park Cave Preserve

2.9.1 Introduction

Table 15. Woodland Park Cave Preserve snapshot showing basic preserve information.

Woodland Park Cave Preserve Snapshot	
Preserve Inception Year	2012
Acreage	10.2
Fence Status	None
Sign Status	Warning signs on all cave gates
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Cat, Duckworth Bat
Non-gated Caves	None

The WCCF agreed to take over management of Duckworth Bat Cave and Cat Cave in 2012 due to a pre-enforcement settlement agreement between the USFWS and Mr. Jimmy Jacobs (President of Shel-Jenn, Incorporated). Shel-Jenn, Incorporated dedicated two conservation areas located in the Woodland Park subdivision, approximately 2.3 miles northwest from Heritage Oaks (Figure 17). The Cat Cave conservation area 4.6 acres and contains the cave's entire surface drainage area (Figure 18). The Duckworth Bat Cave conservation area is 5.6 acres and also contains the cave's entire surface drainage area (Figure 19). Both caves are known to contain the Bone Cave harvestman and both conservation areas contain the majority of the surface troglodite foraging areas proposed by Taylor et al. (2005). Cave gates prevent unauthorized access to Cat Cave and Duckworth Bat Cave while fencing is not currently present.

2.9.2 Hydrogeology

Both conservation areas are located within the EARZ. Based on informal studies conducted for the previous landowner, site-specific geology indicates that these features may be located within the drainage basin for Cowan Spring, which is known to be occupied by the Georgetown salamander (Figure 20).

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan




Figure 17. Woodland Park location map.



Figure 18. Cat Cave conservation area.



Figure 19. Duckworth Bat Cave conservation area.

Hydrogeologic Cross Section of the North San Gabriel River / Cowan Creek Drainage Divide within the Edwards Aquifer Recharge Zone




Figure 20. Cat and Duckworth Bat Caves hydrogeologic setting with local springs.

2.9.3 Caves of the Woodland Park Cave Preserve

Cat Cave and Duckworth Bat Cave maps are included in Appendix F.

Cat Cave: This cave description is paraphrased from Mike Warton and Associates (1999). A cedar elm tree canopy hangs over the entrance to the cave (Photograph 27). The portal is approximately 8 feet wide, rimmed with solid rock, and is located in a surface sink depression that measures approximately 30 feet across. The entrance is in-filled with naturally eroded materials. The entrance room measures approximately 40 feet by 30 feet and gradually slopes to the west. The ceiling height ranges from 1 to 4 feet. Several more passages and rooms exist beyond this point. A gate currently prevents unauthorized access and the Bone Cave harvestman is known from this feature.



Photograph 27. Cat Cave entrance.

Duckworth Bat Cave: This cave description is paraphrased from Mike Warton and Associates (1999). The entrance to the cave is located in a surface sinkhole measuring approximately 10 feet in diameter and is formed along a rock joint trending N20W. The portal is a rock-rimmed vertical shaft approximately 6 feet by 4 feet. After a 27-foot drop to the floor, the room below extends in all directions with ceiling heights between 2 and 6 feet. After excavation, the cave measured at 528.5 feet long and 51.3 feet deep (Photographs 28 and 29). A gate currently prevents unauthorized access and the Bone Cave harvestman is known from this feature.

Table 18 shows documented species within the Woodland Park Cave Preserve.



Photograph 28. Duckworth Bat Cave entrance.



Photograph 29. Duckworth Bat Cave entrance pit.

Table 18. Species previously documented from the Woodland Park Preserve.

Species		Duckworth Bat Cave	Cat Cave
Permitted Species			
<i>Texella reyesi</i>		√	√
Other Species			
Crickets	<i>Ceuthophilus</i> sp.	√	√
Arachnids	<i>Cicurina varians</i>	√	√
	<i>Cicurina vibora</i>		√
	<i>Achaearanea</i> sp.		√
	Araneae	√	√
	<i>Leiobunum</i> sp.	√	
	<i>Pseudouroctonus reddelli</i>	√	√
Millipedes/ Centipedes	<i>Cambala speobia</i>	√	√
	<i>Speodesmus bicornorus</i>	√	
	<i>Scutigera</i> sp.		√
	<i>Scolopendra</i> sp.	√	
Beetles	Staphylinidae	√	
Reptiles/ Amphibians	<i>Lithobates berlandieri</i>	√	√
	<i>Incilius valliceps</i>		√
	<i>Eleutherodactylus marnockii</i>		√
Mammals	<i>Perimyotis subflavus</i>	√	
	<i>Myotis valliceps</i>	√	√
Other	Desert Cockroach		√
	Gnat	√	
	Fly	√	√
	<i>Collembola</i> sp.	√	√
	Flea		√
	Assassin Bug	√	√
	Isopod	√	√
	<i>Helicodiscus eigenmanni</i>	√	

2.10 Karankawa Cave Karst Fauna Area

2.10.1 Introduction

Table 16. Karankawa Cave KFA snapshot showing basic preserve information.

Karankawa Cave KFA Snapshot	
Preserve Inception Year	2012
Acreage	61.7
Fence Status	Northwest boundary fenced along Ronald Reagan Blvd
Sign Status	None
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Pemmican, Angostura
Non-gated Caves	Armadon, Karankawa, Polaris, Quahadi, Snake Dancer, War Party

Williamson County closed on the acquisition of the Karankawa Cave KFA in 2012, as approved by the USFWS. The 61.7-acre Karankawa Cave KFA is located south of SH 195 and approximately 7 miles northwest from the City of Georgetown (Figure 21).

Vegetation within the Karankawa Cave KFA consists of plateau live oak-juniper savanna on gently rolling uplands generally draining westward. A minor upland tributary to Berry Creek runs through the KFA's middle from northeast to southwest and the drainage orientation controlled by a fault that has heavily influenced karst development. The Karankawa Cave KFA boundaries accommodate a biological connection between subsurface mesocaverns and surface flora and fauna within the KFA to the Berry Creek floodplain. The Karankawa Cave KFA is within the northern BFZ and within the Edwards Aquifer's northern segment recharge zone.

The Karankawa Cave KFA contains eight caves (see Figure 21) and two endangered karst invertebrate species, along with at least six "additional species" addressed in the Williamson County RHCP. The Karankawa Cave KFA and associated caves were partially documented and surveyed for biota in 1994 for the Sun City, Georgetown take avoidance plan. Karankawa Cave is known to contain the Bone Cave harvestman and *Batrissodes cryptotexanus*; while War Party, Polaris, and Pemmican Caves are known to contain the Bone Cave harvestman. *Batrissodes cryptotexanus* may yet be detected in War Party Cave; while the presence of other non-listed troglobitic species in Pemmican, Polaris, and Snake Dancer caves indicate significant likelihood of detecting the listed species in those locations.

Fencing currently separates the Karankawa Cave KFA from Ronald Reagan Boulevard, but does not otherwise encircle the entire preserve. Developmental encroachment has continued and the need to separate the eastern and western preserve boundaries has increased. As such, the Karankawa Cave KFA is expected to be fully fenced in the near future. Cave gates currently prevent unauthorized access to Angostura and Pemmican Caves; whereas Armadon, Karankawa, Polaris, Quahadi, Snake Dancer, and War Party Caves are not gated.

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan




Figure 21. Karankawa Cave KFA location map.

2.10.2 Hydrogeology

The Karankawa Cave KFA occupies a relatively flat upland area that drains primarily south and west into an un-named tributary to Berry Creek. Total relief is approximately 40 feet and ranges from approximately 870 to 830 feet amsl. The easement is entirely underlain by Edwards limestone, which is up to 120 feet thick in the area (Collins 2002; Senger et al. 1990).

The Karankawa Cave KFA is within the EARZ northern segment and groundwater recharge primarily occurs in areas where the Edwards Group and upper confining units are exposed at the surface. The tributary location and path appears controlled by the previously mentioned fault. Mapped contact elevations between the Edwards, Comanche Peak and Georgetown formations indicate the fault creates an approximately 15-foot displacement (Figure 22). Cave formation was likely driven by fault-induced, fracture-enhanced permeability. Caves within the Karankawa Cave KFA show fractures related to this fault influence the morphology. Pemmican, Polaris, and Snake Dancer Caves are elongated along fractures that are parallel or sub-parallel to the fault; while Karankawa, War Party, and Angostura caves are influenced by conjugate faults that are roughly perpendicular to the fault. Fracture-driven solutional enlargement has likely enhanced mesocavern connectivity along and across the fault plane.

Rainfall that is not lost through runoff or evapotranspiration likely becomes groundwater by infiltrating through the permeable Edwards limestone. The on-site creek bed is likely an efficient area of focused recharge. Based on the available information, groundwater within the KFA likely flows northeast to southwest mirroring the trend of the caves in the northern cluster. As it approaches the fault the increasing abundance of vertical fractures likely provides numerous avenues for vertical infiltration to the water table which occurs at just above 800 feet, according to nearby well records.




Figure 22. Karankawa Cave KFA hydrogeology.

2.10.3 Caves of the Karankawa Cave Karst Fauna Area

Cave maps are included in Appendix G.

Angostura Cave: The entrance to Angostura Cave drops 4 feet to a ledge-like landing that slopes downward into the cave. The interior is a low, wide irregular room measuring roughly 20 feet by 15 feet by 2 feet high with an uneven floor and nodular ceiling. The floor is soil and clay with breakdown blocks and boulders. The cave seems to have formed by the collapse of a deeper void network and may be a strong indication of mesocavern habitat related to other caves in the northern cluster. A cave gate is currently installed on Angostura Cave.

Armadon Cave: Armadon Cave is a small, shallow cave formed along a bedding plane and measures roughly 10 feet by 15 feet, with at least two humanly inaccessible extensions into the mesocavern matrix and seems to have formed by a deeper void network collapse. This cave is currently not gated.

Karankawa Cave: Karankawa Cave is the largest and deepest cave within the Karankawa Cave KFA and is roughly 92 feet long by 40 feet wide and 21.6 feet deep at its lowest point. Passage proceeds east and then south from the entrance in an arc surrounding a central collapse zone. Karankawa Cave is currently not gated and the Bone Cave harvestman is documented from this location.

Pemmican Cave: Pemmican Cave entrance measures approximately 2 feet long by 1.5 feet wide in solid bedrock, and drops about 10 feet to a floor lined with sediment and small rocks. The main cave extends northeast from the entrance and is a roughly rectangular room measuring approximately 30 feet wide by 50 feet long by up to 4 feet high. An opening to the south becomes too low for human exploration after a few feet. A breakdown-floored alcove in the main room's southeast side gradually becomes too narrow to access. The main room terminates in a narrow passage formed along a joint trending N18E. Near the end of the cave, it is possible to descend through breakdown to enter a low descending lower level passage that extends back towards the entrance for about 25 feet before splitting and becoming too small, which is the deepest cave extent at 17.3 feet below the entrance. Pulverulite is exposed in some places and speleothems are abundant in several areas. Polaris, Pemmican and Snake Dancer Caves are all formed within the fault's damage zone fault and are likely interconnected through mesocaverns. A cave gate is currently installed on Pemmican Cave and the Bone Cave harvestman is documented from this location.

Polaris Cave: Polaris Cave consists of a low, broad chamber formed along a bedding plane with a footprint approximately 50 feet in diameter. Speleothems in the chamber south from the entrance are aligned with fractures parallel to the fault, which passes roughly 100 feet northwest. No cave gate is on this feature, and the Bone Cave harvestman is documented from this location.

Quahadi Cave: Quahadi Cave consists of a low, broad chamber formed along a bedding plane with a footprint measuring approximately 30 feet in diameter. The cave was likely formed by a deeper void network collapse, and there is a strong mesocavern habitat indication related to other caves in the northern cluster. No cave gate is on Quahadi Cave.

Snake Dancer Cave: The entrance to Snake Dancer Cave measures approximately 6 feet wide by 8 feet long by 4.5 feet deep. A 6-foot-wide crawlway leads into an irregular chamber measuring approximately 45 feet long and between 15 to 30 feet wide. The rear cave extent slopes down at about a 45-degree angle to the deepest point at 9 feet below the entrance. Two passages at the end of the room extend for a few feet before becoming too low to continue. One of these passages is formed along a joint trending N18E and investigators detected airflow. Several large breakdown blocks are present in the main room. The

cave contains numerous small speleothems and several massive flowstone deposits. The floor consists largely of black soil and clay. Snake Dancer Cave is not currently gated.

War Party Cave: War Party Cave consists of a low, broad chamber formed along a bedding plane with a footprint measuring approximately 60 feet in diameter. The cave was likely formed by a deeper void network collapse and there is a strong mesocavern habitat indication related to other caves in the northern cluster. The cave drains to a bedrock fracture running perpendicular to the fault to the southeast. The cave's overall morphology is elongated, and the fracture indicates that War Party may have been formed by the collapse of a deeper, fracture-controlled groundwater conduit. That flow would likely have been intercepted by vertical conduits as it approached the fault, which delivered the groundwater to the aquifer base level. Speleothems in the chamber's south end are aligned with fractures parallel to the fault, which passes roughly 100 feet northwest. A cave gate is not currently installed on War Party Cave and the Bone Cave harvestman is documented from this location.

Table 20 shows documented species within the Karankawa Cave KFA.

Table 20. Species previously documented from the Karankawa Cave KFA.

Species		Armadon	Karankawa	Pemmican	Polaris	Quahadi	Snake Dancer	War Party
Permitted Species								
	<i>Batrisesodes cryptotexanus</i>		√					
	<i>Texella reyesi</i>		√	√	√			√
Other Species								
Crickets	<i>Ceuthophilus sp.</i>	√	√	√	√	√	√	√
Arachnids	<i>Cicurina vibora</i>		√	√	√		√	√
	<i>Cicurina varians</i>	√	√	√	√	√	√	√
	<i>Eidmannella sp.</i>		√	√	√			√
	<i>Agynetta llanoensis</i>			√				
	<i>Tayshaneta sp.</i>		√	√				√
	<i>Achaearenea sp.</i>	√		√				
	<i>Cryptachaea porteri</i>					√		
	<i>Leiobunum sp.</i>		√					
	<i>Tartarocreagris sp. (infernalis?)</i>							√
	<i>Pseudouroctonus reddelli</i>			√	√	√		√
Millipedes/ Centipedes	<i>Speodesmus bicornourus</i>		√	√	√		√	√
	<i>Cambala speobia</i>	√	√	√	√	√	√	√
	<i>Oxidus gracilis</i>			√				
	<i>Scolopendra sp.</i>			√			√	√
	<i>Scutigera sp.</i>			√	√	√	√	
Beetles	Coleoptera (Surface Beetles)			√	√		√	
	<i>Batrisesodes uncicornis</i>		√	√			√	√
	Beetle larva			√				
	<i>Tachyini sp.</i>		√	√			√	
	Staphylinidae		√	√	√	√	√	√
	Dermestidae		√					
	<i>Rhadine noctivaga</i>		√	√	√			√

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan

Species		Armadon	Karankawa	Pemmican	Polaris	Quahadi	Snake Dancer	War Party
Reptiles/ Amphibians	<i>Crotalus atrox</i>	√				√		
	<i>Plethodon albagula</i>				√			
	<i>Incilius valliceps</i>							√
	<i>Eleutherodactylus marnockii</i>			√	√		√	√
Mammals	<i>Procyon lotor</i>			√				√
	<i>Didelphis virginiana</i>						√	
	<i>Sylvilagus floridanus</i>				√			
	<i>Perimyotis subflavus</i>	√	√					
	<i>Myotis velifer</i>	√	√				√	
Other	<i>Texoreddellia aquilonalis</i>				√			
	Campodeidae							√
	Desert Cockroach						√	√
	Isopod	√	√				√	
	Silverfish			√				√
	<i>Collembola</i> sp.		√	√	√	√	√	√
	Assassin Bug	√	√	√	√		√	√
	Hemiptera							√
	Diptera			√				
	Gnat	√			√			
	Flea	√						
	Mite		√	√	√	√	√	
	Red Ant				√			
	<i>Helicodiscus eigenmanni</i>		√	√	√			√
	Surface Snail		√		√			
	Moth						√	
	Mosquito		√		√	√		√
	Proscoptera					√		
	Earthworm	√	√	√				

2.11 Coffin Cave Preserve

2.11.1 Introduction

Table 21. Coffin Cave Preserve snapshot showing basic preserve information.

Coffin Cave Preserve Snapshot	
Preserve Inception Year	2014
Acreage	39.4
Fence Status	Fully fenced
Sign Status	Signs posted around perimeter
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Coffin
Non-gated Caves	None

Coffin Cave has long been one of the most notable caves in Central Texas. Written descriptions go back to at least 1958 and the original Texas Cave Survey (Widener 1958). Since 1963—when cave explorers from Southwestern University first mapped Coffin Cave—it has been known as one of the deepest and most extensive caves in Williamson County. The somewhat ominous name of the cave stemmed from the observation that the vertical entrance shaft is roughly coffin-shaped in plan view. Adding to the mystique of the cave is that its location had been lost to the caving community by the time of the listing process for the Travis and Williamson County karst invertebrates in the early 1990s. The location had not been lost to Mr. Marvin Andres, who owned the cave for more than 50 years and kept access generally restricted. Williamson County purchased the cave from Mr. Andres’ heirs in 2014 and established the approximately 39.4-acre preserve (Figure 23).

Fencing is erected across the entire Coffin Cave Preserve and current warning signs aid in trespassing prevention. A steel “A-frame” structure was erected across the Coffin Cave entrance in December 2016 to facilitate ingress/egress for biota monitoring. The cave gate installation will be completed in 2017.

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan




Figure 23. Location map for the Coffin Cave Preserve.

2.11.2 Hydrogeology

Coffin Cave is a highly efficient aquifer recharge feature whose entrance is immediately adjacent to an ephemeral surface drainage channel. Coffin Cave Preserve is mostly atop the Georgetown Formation even though the entire property is within the EARZ. Only the immediate area around cave entrance and along the Coffin Cave Preserve's lowest lying portions (along the drainage channel adjacent to the cave entrance) is the top of the Edwards Formation exposed. Runoff predominates over recharge on the less permeable Georgetown limestone during rain events. Water is effectively routed to the Edwards outcrop where it enters Coffin Cave. The cave's surface drainage basin (approximately 58 acres) is quite large compared to the average Edwards Formation cave. Some of the original drainage basin may have been removed by the quarry immediately west from the Coffin Cave Preserve (see Figure 23). Coffin Cave's drainage basin entirety could not be included within the conservation area due to the basin's large size.

Coffin Cave's other notable hydrological condition relates to its stratigraphic penetration through the Edwards Aquifer's upper half. Surface runoff is delivered almost to the aquifer base level in a matter of minutes during heavy rainfall events. The cave entrance is a solution-enlarged fracture (trending at approximately 125 degrees) measuring 7 feet long and 2 feet wide. The entrance shaft drops approximately 45 feet from its surface expression. Extensive lateral passage continues from the entrance pit's base to the southeast and north before reaching approximately 60 feet deep.

The Edwards limestone contains three distinct zones with regards to hardness and bedrock competency: an upper hard layer, a middle softer layer, and a lower hard layer. The Coffin Cave map produced by ZARA Environmental ([ZARA] 2010) seems to show that the cave morphology evolved in response to this stratified heterogeneity within the Edwards Formation. The profile of the cave shows that the upper approximately 42.6 feet of the cave comprises a vertical shaft formed along a prominent bedrock fracture trending from the northwest to the southeast (Photograph 30). Cave development was restricted to the fracture plane within this zone. Cave morphology is predominantly lateral where cave development followed the softer beds within the Edwards Formation below this zone at between 42.6 feet to 55.7 feet (Figure 24).



Photograph 30. The Coffin Cave entrance is a vertical shaft dropping into the Edwards Formation from the contact with the overlying Georgetown Formation.




Figure 24. Coffin Cave is roughly formed in the Edwards Formation's upper half.

2.11.3 Caves of the Coffin Cave Preserve

Cave map is included within Appendix H.

Coffin Cave: Coffin Cave is the only known cave within the Coffin Cave Preserve (Photograph 31). Coffin Cave is perhaps best known as the source type locality for the endangered Coffin Cave mold beetle (USFWS 1994b). Subsequent taxonomic work has demonstrated that this was a misnomer, however, as the troglobitic mold beetle from this cave is now known to belong to *Batrisodes cryptotexanus*, the first diagnostic specimen of which came from Dragonfly Cave in Sun City Georgetown. Chandler et al. (2009) proposed new common names for both species: Inner Space Caverns mold beetle for *B. texanus* and Dragonfly Cave mold beetle for *B. cryptotexanus*. ZARA (2010) summarized the fauna known from Coffin Cave in their work for the previous cave owner, which is presented in Table 22:

ZARA (2010) also reported collecting a likely Bone Cave harvestman specimen. Therefore, both listed invertebrates covered under the RHCP are known to occur in Coffin Cave.

Table 22. Species previously documented from the Coffin Cave Preserve.

Species		Coffin Cave
Permitted Species		
<i>Batrisodes cryptotexanus</i>		√
<i>Texella reyesi</i>		√
Other Species		
Crickets	<i>Ceuthophilus cunicularis</i>	√
	<i>Ceuthophilus secretus</i>	√
Arachnids	<i>Cicurina vibora</i>	√
	<i>Cicurina varians</i>	√
	<i>Eidmannella pallida</i>	√
Millipedes/ Centipedes	<i>Scolopendra</i> sp.	√
	<i>Cambala speobia</i>	√
Beetles	Coleoptera	√
	Staphylinidae	√
	<i>Ptomaphagus cavernicola</i>	√
Other	<i>Procyon lotor</i>	√
	<i>Pseudosinella violenta</i>	√



Photograph 31. Coffin Cave entrance.

2.12 Beck Commons Preserve

2.12.1 Introduction

Table 17. Beck Commons Preserve snapshot showing basic preserve information.

Beck Commons Preserve Snapshot	
Preserve Inception Year	2014
Acreage	4.2
Fence Status	Fully fenced
Sign Status	Warning signs on all cave gates and on fencing
Baseline Vegetation Survey Date	None
Owner	Williamson County
Gated Caves	Beck Sewer, Beck Trash
Non-gated Caves	None

Written descriptions for Beck Sewer Cave and Beck Trash Cave go back to at least 1958 and the original Texas Cave Survey (Widener 1958). The Beck Sewer Cave name stems from the fact that it was fitted to operate essentially as a septic tank for a ranch house associated with the Beck Ranch operation in 1954. Most land surrounding the Beck Commons Preserve was part of the historic Beck Ranch (Figure 25). Beck Sewer Cave was made accessible to researchers including James Reddell (Texas Memorial Museum) and his mentor Robert W. Mitchell in January 1964. At that time, the researchers made observations as to the cave's general size and orientation; noted guano deposits from a former bat colony; and made a biological collection including blind spiders, harvestmen (almost certainly the Bone Cave harvestman), and beetles. Beck Sewer Cave remains an opportunity for the study of excess nutrient input on troglotic ecosystems. The following quote is included in the Texas Speleological Survey report from that trip:

The cave contains probably the largest population of Rhadine in the State. Literally thousands were observed in all parts of the cave and some were seen eating cricket eggs.


Beck Sewer Cave continued to function as a sewer until around 1994 when local speleologist Mike Warton conducted a trip to the cave and noted the presence of pools of foul water with maggots. WCCF acquired Beck Commons Preserve in 2014.

Notable historic graffiti is located in the dropdown to the lowest room (Photograph 32) and depicts the signature of E.O. Beck from December of 1941—just a few weeks after the Pearl Harbor attacks that brought the United States into World War II. Local lore has it that the caves of Beck Ranch had been explored as potential bomb shelters during the Cold War; perhaps that idea had an earlier origin.

The entirety of this preserve is fenced and appropriate signage has been placed on the fence. A cave gate has already been placed on Beck Sewer Cave and on Beck Trash Cave.



Figure 25. Beck Commons Preserve location map.



Photograph 32. Inscription from Eugene Beck dating to 1941.

2.12.2 Hydrogeology

The Beck Commons Preserve is located within the Edwards Aquifer Recharge Zone and Beck Sewer Cave is formed within the Edwards limestone. Cave extent, morphology, and fracture control indicates it is likely that the Beck Commons Preserve acts to provide recharge despite the surface catchment area not being well defined. Recharge may be suppressed in the northeast corner where fill material 1 to 2 feet deep has been deposited. Beck Sewer Cave allows access to lower levels, with clearly phreatic chambers that are relatively unaffected by the process of collapse and breakdown (Photograph 33). The Beck Sewer Cave's lower level retains an underground stream appearance that has been drained of permanent flow by the aquifer's base level natural lowering. Active recharge can be observed within the cave most prominently in the lower levels.



Photograph 33. Beck Sewer Cave's phreatic passage development in the lower level.

2.12.3 Caves of the Beck Commons Preserve

Cave maps are included within Appendix I.

Several early cave descriptions on the Beck Commons Preserve had the wrong cave map associated with Beck Sewer Cave and the wrong cave name associated with Beck Trash Cave. The map for nearby Beck Ranch Cave (a cave outside the Beck Commons Preserve) had been associated with Beck Sewer Cave and the name "Beck Tin Can Cave" had been applied to Beck Trash Cave. Both sources of confusion were corrected with a renewed mapping effort in 2010; however, no map for Beck Sewer is attached within the appendix of this report.

Beck Trash Cave is within the Beck Commons Preserve, but will not be surveyed for endangered karst biota due to its small size and relative insignificance compared to Beck Sewer Cave. These features are very likely linked, and the Bone Cave harvestmen is known from both caves.

Table 24 shows documented species within the Beck Commons Preserve. Note that no biota survey data are available for Beck Trash Cave.

Table 24. Species previously documented from the Beck Commons KFA.

Species		Beck Sewer
Permitted Species		
<i>Texella reyesi</i>		√
Other Species		
Crickets	<i>Ceuthophilus cunicularis</i>	√
	<i>Ceuthophilus secretus</i>	√
Arachnids	<i>Anapistula</i> sp.	√
	<i>Cicurina vibora</i>	√
	<i>Cicurina varians</i>	√
	Araneae	√
	Mygalomorph (Tarantula)	√
	<i>Achaeearanea</i> sp.	√
	Tick	√
	<i>Tartarocreagris</i> sp. (<i>infernalis</i> ?)	√
Millipedes/ Centipedes	<i>Speodesmus bicornourus</i>	√
	<i>Cambala speobia</i>	√
	<i>Scutigera</i> sp.	√
	Centipede	√
Beetles	<i>Rhadine subterranea</i>	√
	<i>Tachyini</i> sp.	√
	<i>Batrisodes unicomis</i>	√
	Staphylinidae	√
Other	Isopod	√
	<i>Collembola</i> sp.	√
	<i>Tominotus</i> sp.	√
	Surface Snail	√
	<i>Helicodiscus eigenmanni</i>	√
	Assassin Bug	√
	Fly	√
	Gnat	√
	Black Ant	√

2.13 Shaman Cave Karst Fauna Area

2.13.1 Introduction

Shaman Cave KFA Snapshot	
Preserve Inception Year	2016
Acreage	81.7
Fence Status	None
Sign Status	None
Baseline Vegetation Survey Date	None
Owner	WCCF
Gated Caves	Powwow, Shaman
Non-gated Caves	Borgarigmie Pit, Florence No. 18, Haft Shaft, Hatchet, Shawntee Pit, Squaw

The Shaman Cave KFA is located north from Shell Road between Williams Drive (FM 2338) and State Highway 195 near the City of Georgetown (Figure 26). Most of the 81.7-acre Shaman Cave KFA was originally set aside for karst conservation under the take avoidance plan developed by the Sun City Georgetown developers in 1994. The Bone Cave harvestman is documented from both Shaman and Powwow Caves and *B. cryptotexanus* is documented from Shaman Cave; however, the distribution of six smaller caves within the Shaman Cave KFA indicates extensive mesocavern habitat presence between and around the large caves. Other local endemic cave fauna documented from the Shaman Cave KFA include *Cicurina vibora*, *Speodesmus bicornourus*, and *Rhadine noctivaga*; all are additional species addressed in the RHCP (SWCA et al. 2008).

The flora and fauna have begun a successional transition from overgrazed, managed grassland to a predominantly native woodland/grassland mosaic providing suitable habitat for native invertebrates, reptiles and amphibians, and bird species with the cessation of ranching activities more than 20 years ago. The Shaman Cave KFA occurs within the GCWA range and woodland may become suitable for GCWAs over time.




Figure 26. Shaman Cave KFA location map.

2.13.2 Hydrogeology

The Shaman Cave KFA occurs primarily on a broad, flat upland drained primarily to the west, north, and east by a distinctive bend in Berry Creek. Total relief across Shaman Cave KFA is approximately 50 feet ranging from 790 to 740 feet amsl. The Shaman Cave KFA is entirely underlain by the Edwards Limestone. The species-occupied caves occur primarily within the upper 20 feet of the hilltop and most cave entrances are formed between 770 and 780 feet amsl. Most traversable cave passage occurs between 770 and 760 feet amsl (Collins 2005; Senger et al. 1990) (Figure 27). This distribution suggests the presence of a highly developed, stratigraphically controlled cavernous zone of at least 25 acres which likely extends somewhat continuously beneath the hilltop. This karstic horizon within the bedrock likely provides mesocavern habitat between the two endangered species occupied caves and likely beyond.

No mapped faults occur within the Shaman Cave KFA although fractures related to regional faulting are expressed within all caves. A mapped fault passes approximately 0.5 mile northwest of the Shaman Cave KFA (Collins 2005).

All caves within the Shaman Cave KFA have relatively limited surface drainage areas due to their upland terrain location. The endangered species-occupied caves all draw their surface runoff from the hilltop described above. Therefore all surface runoff reaching the cave entrances and footprints originates from within the Shaman Cave KFA.




Figure 27. Geologic cross section of the Shaman Cave KFA.

2.13.3 Caves of the Shaman Cave Karst Fauna Area

The Shaman Cave KFA contains eight caves with two caves known to contain endangered karst invertebrate species, along with at least six “additional species” addressed in the Wilco RHCP. Only Shaman and Powwow Caves are monitored for karst invertebrates. Appendix J includes cave maps for this KFA.

Shaman Cave: Shaman Cave is by far the largest and most biologically diverse cave within the KFA. It has a single passage trending northwest to southeast for approximately 240 feet. The cave is relatively flat in orientation and shallow with the passage majority formed between 5 and 10 feet below the surface. The entrance to Shaman Cave consists of a 1.5-foot-diameter opening in the bedrock which drops approximately 5 feet to the top of a debris cone located in the middle of the caves main passage. The main passage varies from 1.5 to 5 feet high and its width varies from 15 to 25 feet. The cave is approximately 237 feet long and reaches a maximum depth of 14 feet below the surface. Overall, the cave footprint takes the form of a slightly sinuous groundwater conduit with a northwest/southeast trend. Large floor drains accessible to humans at Shaman Cave’s rear indicate connections to a deeper habitat zone. Shaman Cave is gated (Photograph 34) and interior conditions can be quite spacious (Photograph 35). The Bone Cave harvestman and *Batrises cryptotexanus* are both known from this cave.



Photograph 34. Biota survey crew preparing to enter Shaman Cave.



Photograph 35. Biota survey crew (2016) looking for karst invertebrates within Shaman Cave.

Powwow Cave: Powwow Cave is the second largest cave within the Shaman Cave KFA, with a footprint approximately 80 feet long by 30 feet wide. It is essentially a single breakdown chamber occurring between 5 and 15 feet below the surface. The overall cave morphology trends along northwest/southeast in a similar manner to Shaman Cave. The cave contains active speleothems including cave coral, stalactites and stalagmites. The entrance of Powwow Cave is gated (Photograph 36) and occurs in a depression measuring approximately 10 feet long by 4 feet wide. The Bone Cave harvestman is known from this cave.



Photograph 36. Entrance to Powwow Cave.

Additional Caves

The Shaman Cave KFA includes six caves that are not known to contain endangered karst invertebrates. These features are relatively small and shallow for researchers; however, the collective karst feature distribution suggests that Shaman Cave KFA's core is an interconnected cavernous and mesocavernous habitat area spanning more than 25 acres beneath the previously described hilltop. The significant caves' geomorphology suggests they formed from the collapse of a larger underlying karst void network that was likely formed under phreatic conditions in response to paleo-aquifer karst hydrology as groundwater followed an easterly to northeasterly path toward the Edwards Aquifer's confined zone. The current cave floors now are largely formed from breakdown material that has collapsed from the original cave ceiling. The modern cave network has chaotic morphology relative to the paleo cave network because the collapse process is incomplete. A relatively continuous network of voids formed in response to dissolution by flowing groundwater and is now an extensive mesocavern network largely inaccessible to human but readily available to karst invertebrates.

Borgarigmie Pit Cave: The Borgarigmie Pit Cave entrance is an oval opening measuring approximately 3 by 3 feet at the base of a shallow depression. The entrance shaft drops 13 feet to a low room measuring approximately 15 feet long and 7 feet wide. The room is formed along a fracture trending north 85 degrees east. This cave is not gated and is not monitored.

Florence No. 18 Cave: An entrance to Florence No. 18 Cave, measuring roughly 3 feet wide by 4 feet long, drops approximately 8 feet to a slope leading down into an irregular chamber aligned with a southwest-trending fracture. This cave is not gated and is not monitored.

Haft Shaft Cave: The Haft Shaft Cave entrance measures roughly 1.5 feet wide by 2 feet long and drops approximately 13 feet into one end of a low bedding plane room, measuring roughly 12 feet long by 6 feet wide. This cave is not gated and is not monitored.

Hatchet Cave: The entrance to Hatchet Cave is 8 feet wide, with a one-foot-high slope leading to a series of subsequent drops and slopes that reach 19 feet deep, exhibiting an approximately 38-foot passage. This cave is not gated and is not monitored.

Shawntee Pit Cave: The Shawntee Pit Cave is a 12.5-foot-deep pit formed along a north/south-oriented fracture that has been solution-enlarged along an 8-foot-long area. The shaft width is between 1 and 4 feet. A sloping passage extends for another 5 feet from the shaft base. This cave is not gated and is not monitored.

Squaw Cave: Squaw is a single bedding plane room measuring approximately 14 feet long and trending southeast. The cave is formed between 2 and 4 feet below the surface and likely resulted from the collapse of a deeper void. This cave is not gated and is not monitored.

Table 25 shows documented species within the Shaman Cave KFA.

Table 25. Species previously documented from the Shaman Cave KFA.

Species		Shaman	Powwow	Borgarigmie Pit	Florence No. 18	Haft Shaft	Hatchet	Shawntee Pit	Squaw Cave
Permitted Species									
<i>Batrisesodes cryptotexanus</i>		√							
<i>Texella reyesi</i>		√	√						
Other Species									
Crickets	<i>Ceuthophilus sp.</i>	√	√	√	√	√			
	<i>Cicurina vibora</i>	√	√		√				
Arachnids	<i>Cicurina varians</i>	√	√		√		√		
	<i>Anapistula sp.</i>	√							
	<i>Tayshaneta sp.</i>	√	√						
	<i>Achaearanea sp.</i>	√							
	<i>Cryptachaea porteri</i>	√							
	<i>Pseudouroctonus reddelli</i>	√	√						
	<i>Speodesmus bicornourus</i>	√	√				√		
Millipedes/ Centipedes	<i>Cambala speobia</i>	√	√		√		√		
	<i>Rhadine noctivaga</i>		√						
Beetles	<i>Batrisesodes uncicornis</i>		√		√				
	<i>Anillinus affabilis</i>	√							
	Beetle larva	√							
	<i>Crotalus atrox</i>	√							
Reptiles/ Amphibians	<i>Eleutherodactylus marnockii</i>	√							
	Isopod	√							
Other	Assassin Bug	√							
	<i>Texoreddellia aquilonalis</i>	√							
	<i>Collembola sp.</i>	√	√		√		√		
	Gnat		√						
	Mite	√	√	√					

Preserve Descriptions of Land Maintained by the Williamson County Conservation Foundation
under the Williamson County Regional Habitat Conservation Plan

Species	Shaman	Powwow	Borgarigmie Pit	Florence No. 18	Haft Shaft	Hatchet	Shawntee Pit	Squaw Cave
Red Ant	√		√		√		√	√
<i>Pogonomyrmex barbatus</i>	√							
Surface Snail	√	√						
Moth	√	√						
Proscoptera	√							
Earthworm		√						

3.0 LITERATURE CITED

- Chandler, D.S. and J.R. Reddell. 2001. A review of the ant-like litter beetles found in Texas caves (Coleoptera: Staphylinidae: Pselaphinae). *Texas Memorial Museum, Speleological Monographs* 5:115-128.
- Chandler, D.S., J.R. Reddell, and P. Paquin. 2009. New cave Pselaphinae and records from Texas with a discussion of the relationships and distributions of the Texas troglobitic Pselaphinae (Coleoptera: Staphylinidae: Pselaphinae). *Texas Memorial Museum, Speleological Monograph* 7: 125-140.
- Cokendolpher, J.C. 2004. Notes on troglobitic *Cicurina* (Araneae: Dictynidae) from Fort Hood, Texas, with description of another new species. *Texas Memorial Museum, Speleological Monographs* 6:59–62.
- Collins, E.W. 2002. Geologic map of the west half of the Taylor, Texas, 30×60 Minute Quadrangle: central Texas urban corridor, encompassing Round Rock, Georgetown, Salado, Briggs, Liberty Hill, and Leander. University of Texas Bureau of Economic Geology
- . 2005. Geologic Map of the West Half of the Taylor 30×60 Quadrangle: Central Texas Urban Corridor, Encompassing Round Rock, Georgetown, Salado, Briggs, Liberty Hill, and Leander. Bureau of Economic Geology: The University of Texas at Austin. Austin Texas.
- Elliott, W.R. 2004. *Speodesmus* cave millipedes. Four new species from Central Texas (Diplopoda: Polydesmida: Polydesmidae). *Texas Memorial Museum, Speleological Monographs* 6:163-174.
- Horizon Environmental Services, Inc. 2002. Geologic Assessment for 558-acre Williamson County Regional Park, Mayfield Ranch, 3000 CR 175, Round Rock, Williamson County, Texas. Prepared for Williamson County Commissioners Court, Georgetown, TX. January 2002.
- Jones, I.C. 2003. Groundwater availability modeling: Northern segment of the Edwards Aquifer, Texas. Report written for the Texas Water Development Board.
- Krejca, J.K. and B. Weckerly. 2007 Detection Probabilities of Karst Invertebrates. Section 6 Grant Report prepared for the Texas Parks and Wildlife Department
- Mike Warton and Associates. 1999. Karst feature investigations (Phases No. 2 & 3, Project No. 3) lower western side of the “Mayfield/Nelson” Ranch Property located along F.M. Hwy. 1431, Round Rock Area, Williamson County, Texas.
- . 2000. Karst Terrains Survey for Endangered Invertebrate Species Habitat(s) and Associated Point Recharge Value of Karst Features (Phases No. 2 and 3 of Environmental Site Study), 558 Acre Tract Located Along County Road No. 175, Round Rock Area North, Williamson County, Texas. Prepared for Horizon Environmental Services, Inc. August 15, 2000.
- Paquin, P., and M. Hedin. 2004. The power and perils of ‘molecular taxonomy’: a case study of eyeless and endangered *Cicurina* (Aranea: Dictynidae) from Texas caves. *Molecular Ecology* 13(10):3239–3255.

- . 2005. Preliminary results: genetic and morphological analysis of the species limits in *Cicurina* spiders (Aranea: Dictynidae) from southern Travis and northern Hays Counties (TX), with emphasis on *Cicurina cueva* Gertsch and relatives. Unpublished report prepared for the U.S. Fish and Wildlife Service.
- Schneider, K., and D.C. Culver. 2004. Estimating subterranean species richness using intensive sampling and rarefaction curves in a high density cave region in West Virginia. *Journal of Cave and Karst Studies* 66(2): 39–45.
- Senger, R.K., E.W. Collins, and C.W. Kreidler. 1990. Hydrogeology of the Northern Segment of the Edwards Aquifer, Austin Region, Report of Investigations No. 192. Bureau of Economic Geology, University of Texas, Austin.
- Van Kampen-Lewis, S., and K. White. 2017. Management Plan for the Lands Managed by the Williamson County Conservation Foundation under the Williamson County Regional Habitat Conservation Plan. Prepared for the Williamson County Conservation Foundation.
- SWCA Environmental Consultants (SWCA), Smith, Robertson, Elliott, Glen, Klein & Bell, L.L.P., Prime Strategies, Inc., Texas Perspectives, Inc. 2008. Williamson County Regional Habitat Conservation Plan. Prepared for Williamson County Conservation Foundation and The Honorable Lisa Birkman, President and Commissioner, Precinct 1.
- Taylor, S.J., J.K. Krejca, and M.L. Denight. 2005. Foraging range and habitat use of *Ceuthophilus secretus* (Orthoptera: Rhaphidophoridae), a key troglodene in central Texas cave communities. *American Midland Naturalist* 154(1):97–113.
- Ubick, D., and T.S. Briggs. 1992. The harvestman family Phalangodidae. 3. revision of *Texella* Goodnight and Goodnight. *Texas Memorial Museum, Speleological Monographs* 3:155–240.
- U.S. Fish and Wildlife Service (USFWS). 1994a. Recovery plan for endangered karst invertebrates in Travis and Williamson Counties, Texas. Albuquerque, New Mexico.
- . 1994b. 90-day finding on a petition to delist seven Texas karst invertebrates. *59 Federal Register* 11755 11758
- . 2009a. *Bone Cave Harvestman (Texella reyesi) 5-Year Review: Summary and Evaluation*. U.S. Fish and Wildlife Service, Austin Ecological Services Field Office, Austin, Texas.
- . 2009b. *Coffin Cave Mold Beetle (Batrissodes texanus) 5-Year Review: Summary and Evaluation*. U.S. Fish and Wildlife Service, Austin Ecological Services Field Office, Austin, Texas.
- Veni and Associates. 1992. Geologic controls in cave development and the distribution of cave fauna in the Austin, Texas, region. Report for the U.S. Fish and Wildlife Service. Austin, Texas. 77 pp.
- Veni, G. and C. Martinez. 2007. Revision of karst species zones for the Austin, Texas, area. George Veni and Associates. Report for Texas Parks and Wildlife Department. Austin, Texas. 45 pp.
- Verdoorn, R. 1994. Endangered species assessment for Sun City, Georgetown, Georgetown, Texas. Compiled by Richardson Verdoorn for Del Webb Corporation. Austin, Texas.
- White, K. 2009. Hydrologic evolution of the Edwards Aquifer recharge zone (Balcones fault zone) as recorded in the DNA of eyeless *Cicurina* cave spiders, south-central Texas. *Geology* 37(4):339–342.

- White, K, S.W. Carothers, and C. Berkhouse. 2001. The karst fauna region concept and implications for endangered karst invertebrate recovery in Bexar County, Texas. Proceedings of the 2001 National Cave and Karst Management Symposium, Tucson, Arizona.
- Widener, D.L. (ed.). 1958. Texas Cave Survey, Vol 1, No. 1. Reprinted in *Speleo Digest* (1958):1-317.
- ZARA Environmental, LLC (ZARA). 2010. Karst feature investigations on the Katy Whitis property, Williamson County, Texas. ZARA Environmental LLC, Manchaca, Texas. Prepared for Lapeze & Dunahoe, Houston, Texas.

APPENDIX A

Caves of the Twin Springs Karst Fauna Area

Sunless City Cave
Twin Springs Preserve
Williamson County, TX

Tape and Compass Survey by Kemble
White, P.G., and Hub Bechtol, 2000.
Draft by Hub Bechtol.




Figure 5. Map of Sunless City Cave.

Whitney West Cave

Twin Springs Preserve

Williamson County, TX

Tape and Compass Survey by Craig Crawford and Bryan Parker 2009.
Draft by Kemble White.

Plan View



Sinkhole rim

Entrance

North - South Profile View

Entrance

Sinkhole rim

Strong air-flow from
mesocavern matrix

Strong
air-flow

Feet

0 10 20

Figure 6. Map of Whitney West Cave.

APPENDIX B

Cave Maps of the Beck Preserve

BECK CREEK CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY

DATE: NOVEMBER 6, 1991

PERSONNEL:

CHARLY SAVVAS - LEE JAY GRAVES - MIKE WARTON

DRAFT BY: MIKE WARTON

LENGTH: 61.3" DEPTH: 12'.6"

CONFIGURATION OF: 15' X 47'

DIG LEADS

PLAN

0' 1' 2' 3' 4' 5' 10' 15'
SCALE IN FEET: 1" = 5'.0"

PERIMETER EDGE OF
SURFACE SINK

ENTRANCE
(CAVE GATE)
BACFILLED
AREA

ENTRANCE
(CAVE GATE)

PROFILE

SOUTHEAST LOOKING NORTHWEST
ALONG LINE "A - A"

EDWARDS LIMESTONE
MEMBER NO.1
(BASAL UNIT)

DIG
LEADS

12'.6"

T.S.S. 1991

BECK CREVICE CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY

DATE: NOVEMBER 6, 1991

PERSONNEL:


CHARLY SAVVAS - LEE JAY GRAVES - MIKE WARTON

DRAFT BY: MIKE WARTON

LENGTH: 101' 5" DEPTH: 18' 2"

CONFIGURATION OF: 22' 0" X 38' 0"


PLAN



T.S.S. 1991

BECK CREVICE CAVE

FIGURE NO. 2
VERTICAL PROFILE
SOUTH LOOKING NORTH ALONG LINE "A-A"



BECK TEX-2 CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY

DATE: NOVEMBER 6, 1991

PERSONNEL:

CHARLY SAVVAS - LEE JAY GRAVES - MIKE WARTON


DRAFT BY: MIKE WARTON

LENGTH: 69'.6"

DEPTH: 10.6

CONFIGURATION OF: 20' X 32'

PLAN



SCALE IN FEET: 1" = 5'.0"


BECK TEX-2 CAVE

WILLIAMSON COUNTY, TEXAS

FIGURE NO. 2

VERTICAL PROFILE

SOUTH LOOKING NORTH ALONG LINE "A-A"



SCALE IN FEET:

1" = 5' . 0"

BECK SALAMANDER CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTO & TAPE SURVEY

DATE: NOVEMBER 6, 1991

PERSONNEL:

CHARLY SAVVAS-LEE JAY GRAVES - MIKE WARTON

DRAFT BY: MIKE WARTON

LENGTH: 11.4"

DEPTH: 12.1"

PLAN

ENTRANCE
(CAVE GATE)



210°

0' 1' 2' 3' 4' 5' 10' 15'

SCALE IN FEET: 1" = 5'. 0"

ENTRANCE
(CAVE GATE)

AXIAL
(FRACTURE)
TREND

30°

PROFILE
SOUTH LOOKING NORTH



12.1"

EDWARDS LIMESTONE
MEMBER NO.1
(BASAL UNIT)


CLAY FILLED

BECK BAT CAVE

WILLIAMSON COUNTY, TEXAS


SUUNTOS & TAPE SURVEY - DATE: MAY 8, 1996
 PERSONNEL: L. AASE - D. WETUSKI - M. GUERRA - C. WARTON - M. WARTON
 DRAFT BY: M. WARTON
 LENGTH: 182.3' DEPTH: 22.0'
 CONFIGURATION OF: 163' X 91'
 UNITS IN FEET & TENTHS
 ⑥ = CEILING HEIGHTS IN FEET

PLAN



PROFILE

EAST LOOKING WEST
 ALONG AXIAL DEVELOPMENT TREND




BECK PRIDE CAVE

WILLIAMSON COUNTY, TEXAS


SUUNTOS & TAPE SURVEY · DATE: MAY 20, 1996
 PERSONNEL: L. AASE · B. WETUSKI · M. GUERRA · C. WARTON · M. WARTON
 DRAFT BY: M. WARTON
 LENGTH: 407.2' DEPTH: 40.7'
 CONFIGURATION OF: 186' X 72'
 UNITS IN FEET & TENTHS
 (2) = CEILING HEIGHTS IN FEET

PLAN



PROFILE

SOUTHEAST LOOKING NORTHWEST
ALONG AXIAL DEVELOPMENT TREND



- * EDWARDS LIMESTONE FORMATION (Ked)
MEMBER NO. 1 (BASAL UNIT)
- * ENDANGERED INVERTEBRATE SPECIES SITE
- * INTERMEDIATE RANGE POINT RECHARGE FEATURE

BECK HORSE CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY - DATE: MAY 13, 1996

PERSONNEL: L. AASE - B. WETUSKI - M. GUERRA - C. WARTON - M. WARTON

DRAFT BY: M. WARTON



LENGTH: 208.5' DEPTH: 30.0'

CONFIGURATION OF: 157' x 67'

UNITS IN FEET & TENTHS


⑥ = CEILING HEIGHTS IN FEET

PLAN



PROFILE

SOUTHEAST TO NORTHWEST
ALONG AXIAL DEVELOPMENT TREND



- * EDWARDS LIMESTONE FORMATION (Ked)
- MEMBER NO. 1 (BASAL UNIT)
- * ENDANGERED INVERTEBRATE SPECIES SITE
- * INTERMEDIATE RANGE POINT RECHARGE FEATURE

PREPARED BY: MIKE WARTON & ASSOCIATES

T.S.S. 1996

SCALE IN FEET: 1/8" = 1.0"

APPENDIX CC

Cave Maps of the Southwest Williamson County Regional Park

NO PASEO CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JULY 6, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: N.AASE • A.STAPLES •

B.HINDES • C.LANGILLE • M.WARTON • DRAFT BY: M.WARTON

LENGTH: 36.5' DEPTH: 13.5' CONFIGURATION OF: 26' x 11'


(2') = CEILING HEIGHTS IN FEET

PLAN



PROFILE

S. E. to N. W.



PREPARED BY; MIKE WARTON & ASSOCIATES: JULY, 2000

WILLIAMSON COUNTY, TEXAS

PERSONNEL: MIKE WARTON & ASSOCIATES: N. AASE.


A. STAPLES • B. HINDS • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 88.4' DEPTH: 15.5' CONFIGURATION OF: 64' x 38'

(5') = CEILING HEIGHTS IN FEET

PROFILE

Southwest looking Northeast



MONGO CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTO'S & TAPE SURVEY • JUNE 17, 2000

PERSONNEL: MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •

B. HINDS • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 167.8' DEPTH: 29.3' CONFIGURATION OF: 86' x 41'

(5) = CEILING HEIGHTS IN FEET

* EDWARDS LIMESTONE FM. (Ked)

Member No. 1 (Basal Unit)

* Endangered Invertebrate Species Habitat Cave

* "Intermediate" Range Point Recharge Feature

PLAN

0' 5' 10' 15' 20' 25' 30' 35'
Scale in Feet: 1/8" = 1.0'

Perimeter Line of surface sink
(closed depression drainage)

Upslope ↑

N

ENTRANCE
3' Drop

Outline of lower level
9' Drop (pit)

Axial Development trend
at N. 275° W. suggests
Rock Jointing control.

* Floors: dark gray soil &
clay sediments &
scattered rocks/
Reddish moist clays in
Lower Level.

Soil Sediment Filled

Lg. rock slab

PREPARED BY: MIKE WARTON &
ASSOCIATES: JUNE, 2000

PLATE 1 OF 2

MONGO CAVE

WILLIAMSON COUNTY, TEXAS
SUUNTOS & TAPE SURVEY • JUNE 17, 2000

PROFILE

WEST Looking EAST (Mid-Cross Section)

← North
(Upslope)

- * EDWARDS LIMESTONE FM. (Ked)
Member No. 1 (Basal Unit)
- * Endangered Invertebrate Species Habitat Cave
(Bone Cave Harvestman / *Texella Reyesi*)
- * "INTERMEDIATE" Range Point Recharge Feature

ENTRANCE
3' Drop

Collapsed Bedding

Entrance Chamber

Lower Level

—29.3' (Lowest Point)

Moist Reddish clay deposits



PREPARED BY; MIKE WARTON & ASSOCIATES;
JUNE, 2000

Scale in Feet: $1/8" = 1.0'$

PLATE 2 OF 2

ROCK RIDGE CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JUNE 17, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •

B. HINDS • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 103.7' DEPTH: 14.0' CONFIGURATION OF: 80' x 14'

(4') = CEILING HEIGHTS IN FEET

PLAN



PROFILE

South looking North

* EDWARDS Limestone Fm. (Ked)
Member No. 1 (Basal Unit)

* Endangered Invertebrate Species
Habitat Cave

Continues Low

Blockage

Domes

lg. rock slab

Collapse Area

Soil Sediment Floors (dry)

Moist Clay floors along interior
drainage route

Floor Drain
(blocked)

-14' (lowest point)

(filled)

* "Minor" Range Point Recharge Feature

PREPARED BY; MIKE WARTON & ASSOCIATES:
JUNE, 2000

Scale in Feet: 1/8" = 1.0'

TALUS CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JULY 11, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •

B. HINDES • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 67.3' DEPTH: 24.0' CONFIGURATION OF: 43' x 20'


②' = CEILING HEIGHTS IN FEET

PLAN



PROFILE

West looking East



- * **"NON-HABITAT"** Karst Feature
- * **"MINOR TO INTERMEDIATE"** Range POINT RECHARGE Feature/Value
- * Feature's periphery headwall is **"Structurally Stable"**

PREPARED BY; MIKE WARTON & ASSOCIATES:

JULY, 2000

WILCO CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JUNE 29, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •

B. HINDS • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 47' + DEPTH: 15.7' CONFIGURATION OF: 37' x 27' +


(2') = CEILING HEIGHTS IN FEET

PLAN



PROFILE

Rotated to view looking 90° East



CHOYA CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JULY 10, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •

B. HINDES • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 155.8' DEPTH: 8.4' CONFIGURATION OF: 41' x 34'

②' = CEILING HEIGHTS IN FEET

PLAN




* EDWARDS LIMESTONE FM. (Ked)
Member No. 1 (Basal Unit)

* "NON-HABITAT" Karst Feature


* "MINOR" Range Point Recharge
Feature / Value

Floors: Dark Gray Clay soil
sediments w stone cobbles,
Typical throughout.



PROFILE

Southwest looking Northeast



PREPARED BY; MIKE WARTON & ASSOCIATES:
JULY, 2000

SCALE IN FEET:
1/8" = 1.0'

POPPIN ROCKS CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JUNE 15, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •

B. HINDS • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 18.5' DEPTH: 10.0' CONFIGURATION OF: 15'x18'

PLAN


(3') = CEILING HEIGHTS
IN FEET



PROFILE

WEST to EAST

← North



EDWARDS LIMESTONE FM.
(Ked)
Member No. 1 (Basal Unit)

* "NON-HABITAT" Karst Feature
* "MINOR" Range Point
Research Feature

* Massive collapse &
infilling (No air-flow or drains)

Scale in Feet: 1/4" = 1.0'

PREPARED BY; MIKE WARTON & ASSOCIATES:
JUNE, 2000

FENCE & TRAIL CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JULY 17, 2000

PERSONNEL; MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •


B. HINDES • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 20.0' DEPTH: 7.0'


CONFIGURATION OF: 20' x 5'

(2') = CEILING HEIGHTS IN FEET

PLAN



PROFILE



PREPARED BY; MIKE WARTON & ASSOCIATES:
JULY, 2000

SCALE IN FEET: 1/4" = 1.0'


- * "NON-HABITAT" Karst Feature
- * "MINOR" Range Point Recharge Feature

THROUGH-TRIP CAVE

WILLIAMSON COUNTY, TEXAS


SUUNTOS & TAPE SURVEY • AUGUST 2, 2000
 PERSONNEL; MIKE WARTON & ASSOCIATES: A. STAPLES • B. HINDES •
 B. WETUSKI • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON
 LENGTH: 105.9 DEPTH: 16.0' CONFIGURATION OF: 80' x 32'
 ⑦' = CEILING HEIGHTS IN FEET

PLAN



PROFILE

Southeast looking Northwest



- * Potential to Contain Habitat for Endangered Invertebrate Species
- * "Minor" Range Point Recharge Value

PREPARED BY: MIKE WARTON & ASSOCIATES:
 AUGUST, 2000

SCALE IN FEET: 3/32" = 1.0'

LITTLE DEMON CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JULY 27, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: A. STAPLES • B. HINDES •

C. LANGILLE • B. WETUSKI • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 75.6' DEPTH: 19.0' CONFIGURATION OF: 45" x 14"


N (4') = CEILING HEIGHTS IN FEET

PLAN



PROFILE

South looking North



PREPARED BY; MIKE WARTON & ASSOCIATES:
JULY, 2000


MILLENNIUM CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JULY 28, 2000
PERSONNEL; MIKE WARTON & ASSOCIATES: A. STAPLES • B. HINDES •
C. LANGILLE • B. WETUSKI • M. WARTON • DRAFT BY: M. WARTON


PLAN LENGTH: 82.7' DEPTH: 29.0' CONFIGURATION OF: 80' x 35'

(5') = CEILING HEIGHTS IN FEET



PROFILE

South looking North



PREPARED BY; MIKE WARTON & ASSOCIATES: JULY, 2000

KNUCKLE CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JULY 26, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: A. STAPLES •

B. HINDES • C. LANGILLE • B. WETUSKI • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 29.2' DEPTH: 7.3' CONFIGURATION OF: 26 x 11

(2') = CEILING HEIGHTS IN FEET.


PLAN



PROFILE

Southwest looking Northeast

- * "NON HABITAT" Karst Feature
- * "MINOR" Range Point Recharge Feature
- * Cave is "Structurally Stable."



NUEVO CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • JUNE 29, 2000


PERSONNEL; MIKE WARTON & ASSOCIATES: N. AASE • A. STAPLES •

B. HINDS • C. LANGILLE • M. WARTON • DRAFT BY: M. WARTON

LENGTH: 21.3' DEPTH: 5.9' CONFIGURATION OF: 28' x 10'


(2') = CEILING HEIGHTS IN FEET

PLAN



PROFILE

Southwest looking Northeast



* EDWARDS LIMESTONE FM.(Ked)
Member No. 1 (Basal Unit)

* "NON-HABITAT" Karst Feature

* "MINOR" Range Point Recharge Value


PREPARED BY; MIKE WARTON & ASSOCIATES:
JUNE, 2000

SCALE IN FEET:
1/4" = 1.0'

PROSPECTORS CAVE (F-10) WILLIAMSON COUNTY, TEXAS


SUUNTOS & TAPE SURVEY • FEBRUARY 11, 2002
PERSONNEL; MIKE WARTON & ASSOCIATES: M. REYES
M. WARTON • DRAFT BY: M. WARTON
LENGTH: 64.8' DEPTH: 19.0'

PLAN N



PROFILE

South to North



Data:
Edwards limestone fm.
Kainer Unit/Grainstone &
"Kirschberg" level

"Minor" Range Point Recharge Value


Prepared By; Mike Warton & Associates:
FEBRUARY, 2002



VENTURE CAVE (F-11) WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • FEBRUARY 12, 2002
PERSONNEL; MIKE WARTON & ASSOCIATES: M. REYES
M. WARTON DRAFT BY: M. WARTON
LENGTH: 89.3' DEPTH: 21.2'


PLAN



PROFILE

S.E. to N.W.

Entrance




"INTERMEDIATE" Range Point Recharge Value

Prepared By; Mike Warton & Associates:
FEBRUARY, 2002

APPENDIX DD


Cave Maps of the Chaos Preserve




0 1 2 3 4 5
METERS

CHAOS CAVE WILLIAMSON COUNTY, TEXAS


SUUNTO AND TAPE SURVEY
14 APRIL 2000
PETER SPROUSE, JAMES REDDELL
LENGTH: 20 METERS
DEPTH: 3 METERS





UNDER THE FENCE CAVE
WILLIAMSON COUNTY, TEXAS

SUUNTO AND TAPE SURVEY
3 APRIL 2000
LEE JAY GRAVES, NIKI LAKE, PETER SPROUSE
LENGTH: 6 METERS
DEPTH: 3.5 METERS



APPENDIX EE

Cave Maps of the Priscilla's Well Karst Fauna Area


Priscilla's Cave

Williamson County, Texas


Length: 38.3' Depth: 10.5'
Map Drafted by Lindsey Doubleday
5 September 2007

Modified from Mike Warton & Associates, 1994

Plan



Profile



SWCA
ENVIRONMENTAL CONSULTANTS

Figure 5. Priscilla's Cave Map.

Priscilla's Well Cave

Williamson County, Texas

Length: 68.5' Depth: 19.9'

Configuration of: 48' X 19.9'

Map Drafted by Lindsey Doubleday

10 September 2007

Modified from Mike Warton & Associates, 1994

Plan

Rock Joint
Trend at N.275° W.

Entrance

Priscilla's Well

0' 5' 10' 15'

Profile

Entrance
4.5' Drop

Priscilla's Well
15.0' Drop

0'
5'
10'
15'
20'

SWCA
ENVIRONMENTAL CONSULTANTS

Floor 19.9'

Figure 6. Priscilla's Well Cave Map.


APPENDIX FF

Cave Maps of the Woodland Park Cave Preserve

CAT CAVE

WILLIAMSON COUNTY, TEXAS

SUUNTOS & TAPE SURVEY • APRIL 29, 1999
 PERSONNEL; MIKE WARTON & ASSOCIATES: J. KILLIAN •
 R. PRICE • E. BOYD • D. BOYD • M. WARTON • DRAFT BY: M. WARTON
 LENGTH: 543.4' DEPTH: 30.0' CONFIGURATION OF: 233' x 50'




DUCKWORTH BAT CAVE


WILLIAMSON COUNTY, TEXAS

PLAN

SUUNTOS & TAPE SURVEY • APRIL 23, 1999
PERSONNEL; MIKE WARTON & ASSOCIATES: J KILLIAN
E. BOYD • R. PRICE • M. WARTON • DRAFT BY: M. WARTON
LENGTH: 528.5' DEPTH: 51.3' CONFIGURATION OF: 162' x 60'
⑤ = CEILING HEIGHTS IN FEET



- * EDWARDS LIMESTONE FORMATION (Ked)
Member No. 1 (Basal Unit)
- * ENDANGERED INVERTEBRATE SPECIES HABITAT
- * "MINOR" RANGE POINT RECHARGE FEATURE



SCALE IN FEET: 3/32" = 10'

PREPARED BY; MIKE WARTON & ASSOCIATES:
APRIL, 1999

PLATE NO. 1 OF 2

APPENDIX GG

Cave Maps of the Karankawa Cave Karst Fauna Area


ANGOSTURA CAVE

Williamson County, Texas


Suuntos and Tape Survey
4 May 1994
*Doug Allen, L. J. Graves, Dan Love,
Charley Savvas, Mike Warton, Jim Wolff*
Drafted by Mike Warton

Length: 26.5 ft. Depth: 8.1 ft.

PLAN



PROFILE SW looking NE



ARMADON CAVE

Williamson County, Texas

Suuntos and Tape Survey


3 May 1994

D. Allen, L.J. Graves, D. Love, C. Savvas, M. Warton, J. Wolf

Drafted by Mike Warton


Length: 27.5 ft. Depth: 7.5 ft.

PLAN



PROFILE

SW looking NE






KARANKAWA CAVE

Williamson County, Texas

Suuntos and Tape Survey
3 May 1994
D. Allen, L.J. Graves, D. Love, C. Savvas, M. Wrton, J. Wolff
Drafted by Mike Warton

Length: 164.5 ft. Depth: 11.6 ft.


PLAN



PROFILE
SW looking NE



SIDE PROFILE
NW to SE at entrance




PEMMICAN CAVE

Williamson County, Texas

Suuntos and Tape Survey
4 May 1994
*D. Allen, L.J. Graves, D. Love,
C. Savvas, M. Warton, J. Wolff*
Drafted by Mike Warton


Length: 240 ft. Depth: 17.3 ft.

PLAN



PROFILE

west looking east



POLARIS CAVE



Williamson County, Texas

Suuntos and Tape Survey
5 May 1994

D. Allen, L.J. Graves, D. Love, C. Savvas, M. Warton, J. Wolff
Drafted by Mike Warton


Length: 143.7 ft. Depth: 11.8 ft.

PLAN




PROFILE

SW looking NE



QUAHADI CAVE

Williamson County, Texas



SNAKE DANCER CAVE

Williamson County, Texas

Suuntos and Tape Survey

2 May 1994

D. Allen, L. J. Graves, D. Love,


C. Savvas, M. Warton, J. Wolff

Drafted by Mike Warton

Length: 81.3 ft. Depth: 9 ft.

PLAN

ENTRANCE
4.5 ft. drop



PROFILE
east looking west

ENTRANCE
4.5 ft. drop



WAR PARTY CAVE

Williamson County, Texas

Suuntos and Tape Survey


3 May 1994

D. Allen, L.J. Graves, D. Love, C. Savvas, M. Warton, J. Wolfe

Drafted by Mike Warton


Length: 64.5 ft. Depth: 13 ft.

PLAN




PROFILE

SW looking NE



PROFILE

NW looking SE



APPENDIX HH

Cave Map of the Coffin Cave Preserve

APPENDIX I

Cave Map of the Beck Commons Preserve

	AREA	Tc	C25	C100	I25	I100	Q25	Q100
EX. 1 - 11.3 AC	15	0.42	0.49	7.0	8.9	33.2	49.3	
DEV. 1 - 11.3 AC	10	0.80	0.67	8.2	10.4	55.6	78.7	




APPENDIX J

Cave Maps of the Shaman Cave Karst Fauna Area




PLAN



SHAMAN CAVE
 WILLIAMSON COUNTY, TEXAS
 SUUNTOS & TAPE SURVEY • SEPTEMBER 12, 1994
 PERSONNEL: MIKE WARTON & ASSOCIATES:
 LEE JAY GRAVES • DOUG ALLEN • DAN LOVE • JIM WOLFF
 BRENT BARTLETT • JAMES WARTON • MIKE WARTON
 DRAFT BY: MIKE WARTON
 LENGTH: 237.6' DEPTH: 14'
 CONFIGURATION OF: 236' X 45'
 UNITS IN FEET & TENTHS
 BIOLOGY BY: J. REDDELL
 (15) = CEILING HEIGHTS IN FEET

PROFILE

SOUTHWEST LOOKING NORTHEAST



- * EDWARDS LIMESTONE FM. (Ked)
- MEMBER NO. 1 (BASAL UNIT)
- * ENDANGERED INVERTEBRATE SPECIES SITE
- * MINOR POINT RECHARGE SITE

PREPARED BY: MIKE WARTON & ASSOCIATES

T. S. S. 1994


POWWOW CAVE

Williamson County, Texas

Suuntos and Tape Survey
10 September 1994
*D. Allen, B. Bartlett, L. J. Graves, D. Love,
J. Warton, M. Warton, J. Wolff*
Drafted by Mike Warton


Length: 83.2 ft. Depth: 14.9 ft.

PLAN



PROFILE

SW looking NE



BORGARIGMIE PIT CAVE

Williamson County, Texas

Suuntos and Tape Survey


12 September 1994

D. Allen, B. Bartlett, L. J. Graves,
D. Love, J. Warton, M. Warton, J. Wolff


Drafted by Mike Warton

Length: 18.5 ft. Depth: 14 ft.

PLAN




PROFILE



FLORENCE NO. 18 CAVE
Williamson County, Texas

PLAN




Suuntos and Tape Survey
21 April 1994

*D. Allen, L. J. Graves, D. Love,
C. Savvas, M. Warton, J. Wolff*
Drafted by Mike Warton

Length: 52.8 ft. Depth: 16.0 ft.

PROFILE
SW looking NE





HATCHET CAVE

Williamson County, Texas

Suuntos and Tape Survey
21 April 1994
*D. Allen, L. J. Graves, D. Love,
C. Savvas, M. Warton, J. Wolff*
Drafted by Mike Warton

Length: 38.7 ft. Depth: 19 ft.



SQUAW CAVE

Williamson County, Texas

Suuntos and Tape Survey
10 September 1994
*D. Allen, B. Bartlett, L. J. Graves,
D. Love, J. Warton, M. Warton, J. Wolff*
Drafted by Mike Warton

Length: 18.4 ft. Depth: 5.3 ft.

